
pCMV3-C-FLAG Negative Control Vector (C-terminal FLAG-tagged)




Catalog Number: CV012

Physical Map



Vector Name	pCMV3-C-FLAG-NCV
Vector Size	5953bp
Vector Type	Mammalian Expression Vector
Expression Method	Constitutive, Stable / Transient
Promoter	CMV
Antibiotic Resistance	Kanamycin
Selection In Mammalian Cells	Hygromycin
Protein Tag	FLAG
Sequencing Primer	Forward:T7(TAATACGACTCACTATAGGG) Reverse:BGH(TAGAAGGCACAGTCGAGG)

Schematic of pCMV3-C-FLAG-NCV (Negative Control Vector) Multiple Cloning Sites



Physical Introduction

- > Negative control for the pCMV3-C-FLAG clone.
- Vector sequence is the same as pCMV3-C-FLAG, but multiple cloning sites are removed.
- > Designed for mammalian expression, stable or transient.
- >Hygromycin resistance gene for selection of stable cell lines.

Description

Lot : Please refer to the label on the tube

Shipping carrier :

Each tube contains approximately 10 µg of lyophilized plasmid.

Storage :

The lyophilized plasmid can be stored at ambient temperature for three months.

Plasmid Resuspension protocol

- 1. Centrifuge at 5,000 \times g for 5 min.
- 2. Carefully open the tube and add 100 $\mu I~$ of sterile water
- to dissolve the DNA.

3. Close the tube and incubate for 10 minutes at room temperature.

4. Briefly vortex the tube and then do a quick spin to concentrate the liquid at the bottom. Speed is less than $5000 \times g$.

5. Store the plasmid at $\ -20\ ^\circ \mathrm{C}.$

The plasmid is ready for:

- Restriction enzyme digestion
- PCR amplification
- E. coli transformation
- DNA sequencing