

# pCMV3-C-Myc Negative Control Vector (C-terminal Myc-tagged)




## **Physical Map**



Vector Name	pCMV3-C-Myc-NCV
Vector Size	5959bp
Vector Type	Mammalian Expression Vector
Expression Method	Constitutive, Stable / Transient
Promoter	CMV
Antibiotic Resistance	Kanamycin
Selection In Mammalian Cells	Hygromycin
Protein Tag	Мус
Sequencing Primer	Forward:T7(TAATACGACTCACTATAGGG) Reverse:BGH(TAGAAGGCACAGTCGAGG)

# Schematic of pCMV3-C-Myc-NCV (Negative Control Vector) Multiple Cloning Sites





# **Physical Introduction**

- > Negative control for the pCMV3-C-Myc clone.
- Vector sequence is the same as pCMV3-C-Myc, but multiple cloning sites are removed.
- > Designed for mammalian expression, stable or transient.
- >Hygromycin resistance gene for selection of stable cell lines.

### Description

Lot : Please refer to the label on the tube

#### Shipping carrier :

Each tube contains approximately 10 µg of lyophilized plasmid.

#### Storage :

The lyophilized plasmid can be stored at ambient temperature for three months.

### **Plasmid Resuspension protocol**

- 1. Centrifuge at 5,000  $\times$  g for 5 min.
- 2. Carefully open the tube and add 100  $\mu I~$  of sterile water
- to dissolve the DNA.

3. Close the tube and incubate for 10 minutes at room temperature.

4. Briefly vortex the tube and then do a quick spin to concentrate the liquid at the bottom. Speed is less than  $5000 \times g$ .

5. Store the plasmid at -20 °C.

#### The plasmid is ready for:

- Restriction enzyme digestion
- PCR amplification
- E. coli transformation
- DNA sequencing