

Human Interleukin-21 / IL-21 Protein

Catalog Number: GMP-10584-HNAE



Sino Biological
Biological Solution Specialist

General Information

Gene Name Synonym:

CVID11; IL-21; Za11

Protein Construction:

A DNA sequence encoding the mature form of human IL21 isoform 1 (Q9HBE4-1) (Gln23-Ser155) was expressed and purified with an initial Met.

Source: Human

Expression Host: E. coli

QC Testing

Purity: > 95 % as determined by SDS-PAGE

Bio Activity:

Measured by its ability to induced Interferon gamma secretion by human natural killer lymphoma NK-92 cells. The ED_{50} for this effect is 0.4-2 ng/mL.

Endotoxin:

< 0.01 EU per μ g of the protein as determined by the LAL method

Stability:

Samples are stable for up to twelve months from date of receipt at -70 °C

Predicted N terminal: Met

Molecular Mass:

The recombinant human IL-21 consists of 134 amino acids and has a calculated molecular mass of 15.6 kDa.

Formulation:

Lyophilized from sterile PBS, pH 7.4.

Normally 5 % - 8 % trehalose, mannitol and 0.01% Tween80 are added as protectants before lyophilization. Specific concentrations are included in the hardcopy of COA. Please contact us for any concerns or special requirements.

Usage Guide

Storage:


Store it under sterile conditions at -20°C to -80°C upon receiving. Recommend to aliquot the protein into smaller quantities for optimal storage.

Avoid repeated freeze-thaw cycles.

Reconstitution:

Detailed reconstitution instructions are sent along with the products.

SDS-PAGE:



Protein Description

IL21 belongs to the IL-15/IL-21 family. It is a cytokine with immunoregulatory activity. Cytokines are proteinaceous signaling compounds that are major mediators of the immune response. They control many different cellular functions including proliferation, differentiation and cell survival/apoptosis but are also involved in several pathophysiological processes including viral infections and autoimmune diseases. Cytokines are synthesized under various stimuli by a variety of cells of both the innate (monocytes, macrophages, dendritic cells) and adaptive (T- and B-cells) immune systems. IL21 is expressed in activated CD4-positive T-cells but not in CD8-positive T-cells, B-cells, or monocytes. It may promote the transition between innate and adaptive immunity. IL-21 has been tried as therapy for alleviating allergic responses. It can significantly decrease pro-inflammatory cytokines produced by T cells in addition to decreasing IgE levels in a mouse model for rhinitis (nasal passage inflammation)

References

- 1.Coquet JM, et al. (2007) IL-21 is produced by NKT cells and modulates NKT cell activation and cytokine production. J Immunol. 178(5):2827-34.
- 2.Wei L, et al. (2007) IL-21 is produced by Th17 cells and drives IL-17 production in a STAT3-dependent manner. J Biol Chem. 282(48):34605-10.
- 3.Parrish-Novak J, et al. (2002) Interleukin-21 and the IL-21 receptor: novel effectors of NK and T cell responses. J Leukoc Biol. 72(5):856-63.
- 4.Kuchen S, et al. (2007) Essential role of IL-21 in B cell activation, expansion, and plasma cell generation during CD4+ T cell-B cell collaboration. J Immunol. 179(9):5886-96.

Manufactured By Sino Biological Inc., FOR RESEARCH USE ONLY. NOT FOR USE IN HUMANS.

For US Customer: Fax: 267-657-0217

● Tel: 215-583-7898

Global Customer: Fax :+86-10-5862-8288

● Tel:+86-400-890-9989 ●

● <http://www.sinobiological.com>