

Human CD147 / BSG transcript variant 2 ORF mammalian expression plasmid, N-HA tag



Sino Biological Inc.

Biological Solution Specialist

Catalog Number: HG10186-NY

General Information

Gene :	basigin (Ok blood group) (BSG), transcript variant 2
Official Symbol :	BSG
Synonym :	BSG, M6, OK, 5F7, TCSF, CD147, EMMPRIN
Source :	Human
cDNA Size:	810bp
RefSeq :	NM_198589.1
Plasmid:	pCMV3-HA-BSG

Description

Lot : Please refer to the label on the tube

Sequence Description :

Identical with the Gene Bank Ref. ID sequence.

Restriction site: KpnI + XbaI (6kb + 0.84kb)

Vector : pCMV3-SP-N-HA

Shipping carrier :

Each tube contains approximately 10 µg of lyophilized plasmid.

Storage :

The lyophilized plasmid can be stored at ambient temperature for three months.

Quality control :

The plasmid is confirmed by full-length sequencing with primers in the sequencing primer list.

Sequencing primer list :

pCMV3-F:	5' CAGGTGTCCACTCCAGGTCCAAG 3'
pcDNA3-R :	5' GGCAACTAGAAGGCACAGTCGAGG 3'
Or	
Forward T7 :	5' TAATACGACTCACTATAGGG 3'
ReverseBGH :	5' TAGAAGGCACAGTCGAGG 3'

pCMV3-F and pcDNA3-R are designed by Sino Biological Inc. Customers can order the primer pair from any oligonucleotide supplier.

Plasmid Resuspension protocol

1. Centrifuge at 5,000×g for 5 min.
2. Carefully open the tube and add 100 µl of sterile water to dissolve the DNA.
3. Close the tube and incubate for 10 minutes at room temperature.
4. Briefly vortex the tube and then do a quick spin to concentrate the liquid at the bottom. Speed is less than 5000×g.
5. Store the plasmid at -20 °C.

The plasmid is ready for:

- Restriction enzyme digestion
- PCR amplification
- *E. coli* transformation
- DNA sequencing

E. coli strains for transformation (recommended but not limited)

Most commercially available competent cells are appropriate for the plasmid, e.g. TOP10, DH5α and TOP10F'.

Manufactured By Sino Biological Inc., FOR RESEARCH USE ONLY. NOT FOR USE IN HUMANS.

Fax :+86-10-51029969

• Tel: +86- 400-890-9989

• <http://www.sinobiological.com>

Human CD147 / BSG transcript variant 2 ORF mammalian expression plasmid, N-HA tag



Sino Biological Inc.

Biological Solution Specialist

Catalog Number: HG10186-NY


Vector Information

All of the pCMV vectors are designed for high-level stable and transient expression in mammalian hosts. High-level stable and non-replicative transient expression can be carried out in most mammalian cells. The vectors contain the following elements:

- Human enhanced cytomegalovirus immediate-early (CMV) promoter for high-level expression in a wide range of mammalian cells.
- Hygromycin resistance gene for selection of mammalian cell lines.
- A Kozak consensus sequence to enhance mammalian expression.

Vector Name	pCMV3-SP-N-HA
Vector Size	6146bp
Vector Type	Mammalian Expression Vector
Expression Method	Constitutive, Stable / Transient
Promoter	CMV
Antibiotic Resistance	Kanamycin
Selection In Mammalian Cells	Hygromycin
Protein Tag	HA

Physical Map of Plasmid :



Manufactured By Sino Biological Inc., FOR RESEARCH USE ONLY. NOT FOR USE IN HUMANS.

Fax :+86-10-51029969

• Tel:+86- 400-890-9989

• <http://www.sinobiological.com>

pCMV3-SP-N-HA (suitable for secretory and membrane protein expression)



Sino Biological Inc.
Biological Solution Specialist

Physical Map



Comments for pCMV3-SP-N-HA:

CMV promoter: bases 250-837
enhancer: bases 838-1445
SV40 early promoter: bases 2387-2756
Hygromycin ORF: bases 2774-3799
pUC origin: bases 4442-5115
Kanamycin ORF: bases 5189-6004

Description

Vector Name	pCMV3-SP-N-HA
Vector Size	6146bp
Vector Type	Mammalian Expression Vector
Expression Method	Constitutive, Stable / Transient
Promoter	CMV
Antibiotic Resistance	Kanamycin
Selection In Mammalian Cells	Hygromycin
Protein Tag	HA
Sequencing Primer	Forward:T7(TAATACGACTCACTATAGGG) Reverse:BGH(TAGAAGGCACAGTCGAGG)

Schematic of pCMV3-SP-N-HA Multiple Cloning Sites



pCMV3-SP-N-HA is recommended for constructing the N-HA tag secretory and membrane proteins expression vector which containing a naïve signal peptide. An universal signal peptide is used to instead the naïve signal peptide.