# Human c-Abl / ABL1 Gene ORF cDNA clone in cloning vector

Catalog Number: HG11199-M



# **General Information**

Gene: c-abl oncogene 1, receptor tyrosine

kinase

Official Symbol: ABL1

**Synonym:** ABL; bcr/abl; c-ABL; c-ABL1; JTK7; p150;

v-abl

Source: Human

cDNA Size: 3450bp

**RefSeq:** BC117451.1

Plasmid: pMD-ABL1

# **Description**

Lot: Please refer to the label on the tube

# **Sequence Description:**

Identical with the Gene Bank Ref. ID sequence except for four point mutations: 9 G/A, 12 G/A, 1554 A/G and 3381 A/G not causing the amino acid variation.

Vector:

pMD18-T Simple

#### **Quality control:**

The plasmid is confirmed by full-length sequencing with primers in the sequencing primer list.

#### Sequencing primer list:

M13-47: 5' GCCAGGGTTTTCCCAGTCACGAC 3'

RV-M: 5' GAGCGGATAACAATTTCACACAGG 3'

Other M13 primers can also be used as sequencing primers.

### Shipping carrier:

Each tube contains approximately 10 µg of lyophilized plasmid.

#### Storage:

The lyophilized plasmid can be stored at ambient temperature for three months.

# **Plasmid Resuspension protocol**

- 1. Centrifuge at 5,000×g for 5 min.
- 2. Carefully open the tube and add 100  $\mu l$  of sterile water to dissolve the DNA.
- Close the tube and incubate for 10 minutes at room temperature.
- 4. Briefly vortex the tube and then do a quick spin to concentrate the liquid at the bottom. Speed is less than 5000 $\times$ g.
- 5. Store the plasmid at  $-20 \,^{\circ}$ C.

### The plasmid is ready for:

- Restriction enzyme digestion
- PCR amplification
- E. coli transformation
- · DNA sequencing

# *E.coli* strains for transformation (recommended but not limited)

Most commercially available competent cells are appropriate for the plasmid, e.g. TOP10, DH5 $\alpha$  and TOP10F $^{\prime}$ .

Website: http://www.sinobiological.com

# Human c-Abl / ABL1 Gene ORF cDNA clone in cloning vector

Catalog Number: HG11199-M




# **Vector Information**

pMD18-T Simple Vector is a high-efficiency TA cloning vector constructed from pUC18, of which the initial multiple cloning sites (MCS) were destroyed. The pMD18-T Simple Vector is 2.6kb in size and contains the ampicillin resistance gene for selection. The coding sequence was inserted by TA cloning at site 425.

Notes: The direction of cDNA insertion into the TA-cloning vector is random, maybe forward or reverse. For insert orientation information, please feel free to contact us.

#### Physical Map of pMD18-T Simple (MCS destroyed):



Sequencing Primer M13-47

5'-CGCCAGGGTTTTCCCAGTCACGACGTTGTAAAACGACGGCCAGTGCC

EcoR V

AAAGAAGCATGACGGCAAGTGGACGATATCTCCAGAAGGATCGCCGGGAA

Cloning Site:425
...gtggacgatT atctccaga...
...cacctgcta Ttagaggtct... (TA cloning Site)

CCGAGGACGAGTTCGTAATCATGGTCATAGCTGTTT<u>CCTGTGTGAAATTGTT</u>
Sequencing Primer RV-M

ATCCGCTC -3'

Website: http://www.sinobiological.com