Human IDH2 Gene cDNA clone plasmid

Catalog Number: HG14526-G



General Information

Gene: isocitrate dehydrogenase 2 (NADP+),

mitochondrial

Official Symbol: IDH₂

Synonym: IDH, IDP, IDHM, IDPM, ICD-M, D2HGA2,

mNADP-IDH

Source: Human

cDNA Size: 1359bp

RefSeq: BC009244

Plasmid: pGEM-IDH2

Description

Lot: Please refer to the label on the tube

Sequence Description:

Identical with the Gene Bank Ref. ID sequence.

Vector:

pGEM-T

Shipping carrier:

Each tube contains approximately 10 µg of lyophilized plasmid.

Storage:

The lyophilized plasmid can be stored at ambient temperature for three months.

Quality control:

The plasmid is confirmed by full-length sequencing with primers in the sequencing primer list.

Sequencing primer list:

5' GCCAGGGTTTTCCCAGTCACGAC 3' M13-47:

RV-M: 5' GAGCGGATAACAATTTCACACAGG 3'

Other M13 primers can also be used as sequencing primers.

Plasmid Resuspension protocol

- 1. Centrifuge at 5,000 × g for 5 min.
- 2.Carefully open the tube and add 100 µl of sterile water to dissolve the DNA.
- 3. Close the tube and incubate for 10 minutes at room temperature.
- 4. Briefly vortex the tube and then do a quick spin to concentrate the liquid at the bottom. Speed is less than 5000×g.
- 5.Store the plasmid at -20 °C.

The plasmid is ready for:

- · Restriction enzyme digestion
- · PCR amplification
- · E. coli transformation
- DNA sequencing

E.coli strains for transformation (recommended but not limited)

Most commercially available competent cells are appropriate for the plasmid, e.g. TOP10, DH5α and TOP10F'.

Fax:+86-10-51029969

● Tel:+86- 400-890-9989 ● http://www.sinobiological.com

Human IDH2 Gene cDNA clone plasmid

Catalog Number: HG14526-G



Vector Information

The pGEM-T vector is a high-efficiency TA cloning vector which contains multiple cloning sites as shown below. The pGEM-T vector is 3.0kb in size and contains the amplicin resistance gene for selection. The coding sequence was inserted by TA cloning.

Physical Map of pGEM-T:

