



Catalog Number: HG14693-G

General Information

Gene : hydroxysteroid (17-beta) dehydrogenase 8
Official Symbol : HSD17B8
Synonym : KE6, FABG, HKE6, FABGL, RING2, H2-KE6, SDR30C1, D6S2245E, dJ1033B10.9
Source : Human
cDNA Size: 786bp
RefSeq : BC008185
Plasmid: pGEM-HSD17B8

Description

Lot : Please refer to the label on the tube

Sequence Description :

Identical with the Gene Bank Ref. ID sequence.

Vector :

pGEM-T

Shipping carrier :

Each tube contains approximately 10 µg of lyophilized plasmid.

Storage :

The lyophilized plasmid can be stored at ambient temperature for three months.

Quality control :

The plasmid is confirmed by full-length sequencing with primers in the sequencing primer list.

Sequencing primer list :

M13-47 : 5' GCCAGGGTTTCCCAGTCACGAC 3'

RV-M : 5' GAGCGGATAACAATTTACACAGG 3'

Other M13 primers can also be used as sequencing primers.

Plasmid Resuspension protocol

1. Centrifuge at 5,000×g for 5 min.
2. Carefully open the tube and add 100 µl of sterile water to dissolve the DNA.
3. Close the tube and incubate for 10 minutes at room temperature.
4. Briefly vortex the tube and then do a quick spin to concentrate the liquid at the bottom. Speed is less than 5000×g.
5. Store the plasmid at -20 °C.

The plasmid is ready for:

- Restriction enzyme digestion
- PCR amplification
- *E. coli* transformation
- DNA sequencing

E. coli strains for transformation (recommended but not limited)


Most commercially available competent cells are appropriate for the plasmid, e.g. TOP10, DH5α and TOP10F'.

Catalog Number: HG14693-G

Vector Information

The pGEM-T vector is a high-efficiency TA cloning vector which contains multiple cloning sites as shown below. The pGEM-T vector is 3.0kb in size and contains the ampicin resistance gene for selection. The coding sequence was inserted by TA cloning.

Physical Map of pGEM-T :



pGEM-T Vector

