



Sino Biological Inc.
Biological Solution Specialist

Immunoprecipitation/IP Kit- Anti-NFkBIB / IkB-beta Immunomagnetic Beads

Catalog Number: MB12685-T20

Please read this instruction manual carefully before using the product

Product Contents

Contents	Package 1	Package 2	Storage
Anti-NFKBIB / IκB-beta Immunomagnetic Beads ^{1 3}	1 mL	5 mL	2-8℃ for 12 months
NP40 Cell Lysis Buffer	4 mL	22 mL	-20℃ for 12 months
5×TBST (pH7.4)	Required but not supplied		
1×TBST (pH7.4)	Required but not supplied		
ddH ₂ O	Required but not supplied		
Alkaline Elution Buffer	3 mL	15 mL	2-8℃ for 12 months
Acidity Elution Buffer	3 mL	15 mL	2-8℃ for 12 months
Neutralization Buffer	2 mL	8 mL	2-8℃ for 12 months
Magnetic Separator	Not included (refer related product MAGS001)	One MAGS001 included in China ²	

- [1] The IP KIT contains anti-NFKBIB / IκB-beta Immunomagnetic Beads(2 mg/mL) in phosphate buffered saline (PBS, pH 7.4) with sodium azide (0.1%).
- [2] The Magnetic Separator cannot be included for oversea customers due to shipment prohibition.
- [3] Immunomagnetic Beads kits are shipped at ambient temperature in which immunomagnetic beads are provided in liquid buffer.

Product Description

The Anti-NFKBIB / IκB-beta Immunomagnetic Beads, conjugated with Anti-NFKBIB / IκB-beta antibody, are used for immunoprecipitation (IP) of NFKBIB / IκB-beta proteins which expressed in vitro expression systems and bacterial and mammalian cell lysates.

For IP, the beads are added to a sample containing NFKBIB / IκB-beta proteins to form a bead-protein complex. The complex is removed from the solution manually using a Magnetic Separator. The bound NFKBIB / IκB-beta proteins are dissociated from the Immunomagnetic Beads using an elution buffer.

Antibody Information

- Antibody:** NFKBIB / IκB-beta Antibody, Rabbit PAb, Antigen Affinity Purified(12685-T20)
- Immunogen:** Recombinant Human NFKBIB / IκB-beta Protein
- Clone ID:**
- Isotype:** Rabbit IgG
- Specificity:** Human NFKBIB / IκB-beta
- Guaranteed Applications:** IP, Minimum Protein Purification
- Preparation:** Produced in rabbits immunized with purified, recombinant Human NFKBIB / IκB-beta (rh NFKBIB / IκB-beta; AAH15528.1; 1M-356V). NFKBIB / IκB-beta specific IgG was purified by Human NFKBIB / IκB-beta affinity chromatography.




Fig. 1 Immunoprecipitation (IP) Protocol

Protocol

The protocol (Fig. 1) uses 50 μL Anti-NFKBIB / I κ B-
betaImmunomagnetic Beads, but this can be scaled up or
down as required.

Cell Lysis

Cells may be lysed using any standard cell lysis protocol in
accordance with your starting materials. We suggest using
NP40 Cell Lysis Buffer (supplied with kit).

Immunoprecipitate Target Antigen

1. Add 50 μL of Immunomagnetic Beads into a 1.5 mL
microcentrifuge tube.
2. Add 150 μL of 1 $\times$ TBST buffer to the Immunomagnetic
Beads and gently vortex to mix.
3. Place the tube into a Magnetic Separator to collect the
beads against the wall side of the tube. Remove and discard
the supernatant.
4. Add 1 mL of 1 $\times$ TBST buffer to the tube. Invert the tube
several times or gently vortex to mix for 1 min. Collect
Immunomagnetic Beads with a Magnetic Separator.
Remove and discard the supernatant.
5. Add the sample containing target protein (~100 μg of
protein in 100 μL) to the pre-washed Immunomagnetic
Beads, add 400 μL of 1 $\times$ TBST buffer and incubate at room
temperature for 30 min with mixing.
6. Collect the Immunomagnetic Beads with a Magnetic
Separator, remove the unbounded sample and save for
analysis.
7. Add 300 μL of 5 $\times$ TBST buffer to the tube and gently mix.
Collect the Immunomagnetic Beads and discard the
supernatant. Repeat this wash twice.
8. Add 300 μL of ddH₂O to the tube and gently mix. Collect
the Immunomagnetic Beads on a Magnetic Separator and
discard the supernatant.

Elute Target Antigen.

A. Alkaline Elution Protocols

1. Add 100 μL of Alkaline Elution buffer to the tube.
2. Gently vortex to mix and incubate the sample at room
temperature on a rotator for 5 min.
3. Magnetically separate the Immunomagnetic Beads and
save the supernatant containing the target antigen.
4. To neutralize the sample, add 50 μL of Neutralization
Buffer for each 100 μL of eluate.

B. Acidity Elution

1. Add 100 μL Acidity Elution Buffer.
2. Gently vortex to mix and incubate the sample at room
temperature on a rotator for 5-10 min.
3. Magnetically separate the Immunomagnetic Beads and
save the supernatant containing the target antigen.

4. To neutralize the low pH, add 15 μL of Neutralization
Buffer for each 100 μL of eluate.

C. Elution Using Sample Buffer

1. Add 100 μL of SDS-PAGE sample buffer to the tube.
2. Gently vortex to mix and incubate the sample at 95-
100°C for 5-10 min.
3. Magnetically separate the Immunomagnetic Beads and
save the supernatant containing the antigen.

Reference Information

Related Products

Products	Cat No.
Magnetic Separator-1.5 (2 tubes)	MAGS001
Immunoprecipitation Kit -Immunomagnetic Beads Protein A Kit	BA10600
Immunoprecipitation Kit -Immunomagnetic Beads Protein G Kit	BG13103
Immunoprecipitation Kit -Immunomagnetic Beads Protein L Kit	BL11044
Immunoprecipitation Kit -Anti-DYKDDDDK(Flag®) Tag Immunomagnetic Beads Kit	TB101274
Immunoprecipitation Kit -Anti-GFP Tag Immunomagnetic Beads Kit	TB13105
Immunoprecipitation Kit -Anti-Myc Tag Immunomagnetic Beads Kit	TB100029
Immunoprecipitation Kit -Anti-HA Tag Immunomagnetic Beads Kit	TB100028
Immunoprecipitation Kit -Anti-V5 Tag Immunomagnetic Beads Kit	TB100378

Trouble Shooting

Problem	Possible Cause	Solution
Little or no protein is detected	Protein degraded	Include protease inhibitors (PMSF) in the lysis buffer
		Use new lysate or lysate stored at -80° C
	No or minimal protein was expressed	Verify protein expression by SDS-PAGE or Western blot
		Analysis of the lysate using an positive control as a reference
		Increase the amount of lysate used for IP/Co-IP
		Use a more sensitive detection system

Problem	Possible Cause	Solution
Magnetic Beads aggregated	Magnetic Beads were frozen or centrifuged	Handle the Beads as directed in the instructions
	Buffer was incompatible with magnetic beads	
	Detergent was not added to the wash and bind solutions	
Failure to co-IP interacting protein	Wash conditions were too stringent for the weak or transient interaction	Reduce the number of washes
		Lower the ionic strength of the wash buffer
	Interacting protein was expressed at a low level	Apply additional protein sample
		Use a more sensitive detection system
	Buffer system was not optimal for the protein: protein interaction	Optimize the co-IP buffer
	Insufficient sample was loaded on the gel for Western blot detection	Elute sample in 30% acetonitrile 0.5% formic acid, then
		Bring the sample back up in SDS-PAGE Sample Buffer and load entire elution fraction on