

Human IFN α 4 / IFN α 4 / Interferon alpha-4 Protein (Fc Tag)

Catalog Number: 10336-H01H



Sino Biological
Biological Solution Specialist

General Information

Gene Name Synonym:

IFN-alpha4a; INF4; MGC142200

Protein Construction:

A DNA sequence encoding the mature form of human IFN- α 4a (NP_066546.1) (Cys 24-Asp 189) was expressed with the N-terminal fused Fc region of human IgG1.

Source: Human

Expression Host: HEK293 Cells

QC Testing

Purity: > 97 % as determined by SDS-PAGE

Bio Activity:

Measured in antiviral assays using WISH human amnion cells infected with vesicular stomatitis virus (VSV). The EC₅₀ for this effect is 0.2-1 ng/mL.

Endotoxin:

< 1.0 EU per μ g of the protein as determined by the LAL method

Stability:

Samples are stable for up to twelve months from date of receipt at -70 °C

Predicted N terminal: Glu

Molecular Mass:

The recombinant human Fc/IFN- α 4a is a disulfide-linked homodimeric protein. The reduced monomer consists of 403 amino acids and has a predicted molecular mass of 46 kDa. As a result of glycosylation, the apparent molecular mass of rh Fc/IFN- α 4a monomer is approximately 50 kDa in SDS-PAGE under reducing conditions.

Formulation:

Lyophilized from sterile PBS, pH 7.4

Normally 5 % - 8 % trehalose, mannitol and 0.01% Tween80 are added as protectants before lyophilization. Specific concentrations are included in the hardcopy of COA. Please contact us for any concerns or special requirements.

Usage Guide

Storage:


Store it under sterile conditions at -20°C to -80°C upon receiving. Recommend to aliquot the protein into smaller quantities for optimal storage.

Avoid repeated freeze-thaw cycles.

Reconstitution:

Detailed reconstitution instructions are sent along with the products.

SDS-PAGE:



Protein Description

Interferon, alpha 4 (IFNA4) belongs to the alpha/beta interferon family. Two variants of IFNA4 (IFNA4a and IFNA4b) are known, which differ from each other by changes in their coding regions at nucleotide positions 220 and 410 and can be distinguished by selective restriction enzyme analysis. Interferons are produced by macrophages, IFN-alpha have antiviral activities. Interferon stimulates the production of two enzymes: a protein kinase and an oligoadenylate synthetase. IFN-alpha, the first cytokine to be produced by recombinant DNA technology, has emerged as an important regulator of growth and differentiation, affecting cellular communication and signal transduction pathways as well as immunological control. Originally discovered as an antiviral substance, the efficacy of IFN-alpha in malignant, viral, immunological, angiogenic, inflammatory, and fibrotic diseases suggests a spectrum of interrelated pathophysiologies. IFN-alpha emerged as a prototypic tumor suppressor protein that represses the clinical tumorigenic phenotype in some malignancies capable of differentiation.

References

1. Lau JY, *et al.* (1993) Discrepancy between biochemical and virological responses to interferon-alpha in chronic hepatitis C. *Lancet*. 342(8881): 1208-9.
2. Kessler DS, *et al.* (1990) Interferon-alpha regulates nuclear translocation and DNA-binding affinity of ISGF3, a multimeric transcriptional activator. *Genes Dev.* 4(10): 1753-65.
3. Gutterman JU. Cytokine therapeutics: lessons from interferon alpha. *Proc Natl Acad Sci U S A.* 91(4): 1198-205.

Manufactured By Sino Biological Inc., FOR RESEARCH USE ONLY. NOT FOR USE IN HUMANS.

For US Customer: Fax: 267-657-0217 • Tel: 215-583-7898

Global Customer: Fax :+86-10-5862-8288 • Tel:+86-400-890-9989 • <http://www.sinobiological.com>