

TNFR2 / CD120b / TNFRSF1B Neutralizing Antibody



Catalog Number: 10417-R00N6

EliteRmab® is a registered trademark of Sino Biological Inc.

General Information	
Immunogen:	Recombinant Human TNFR2 / CD120b / TNFRSF1B Protein (Catalog#10417-H08H)
Clone ID:	R006
Ig Type:	Rabbit IgG
Applications:	Neutralization
Specificity:	Human TNFR2 / CD120b / TNFRSF1B
Formulation:	0.2 µm filtered solution in Histidine and Arginine buffer containing 120mM NaCl, 0.02% Tween 80, pH6.0
Storage:	< -20°C

Preparation

This antibody was obtained from a rabbit immunized with purified, recombinant Human TNFR2 / CD120b / TNFRSF1B (rh TNFR2 / CD120b / TNFRSF1B; Catalog#10417-H08H; NP_001057.1; Met1-Asp257) and was produced using recombinant antibody technology.

Specificity

Human TNFR2 / CD120b / TNFRSF1B

No cross-reactivity with Mouse TNFR2 (Catalog#50128-M08H) in ELISA assay

Storage

This antibody can be stored at 2°C-8°C for one month without detectable loss of activity. Antibody products are stable for twelve months from date of receipt when stored at -20°C to -80°C. **Preservative-Free**.

Sodium azide is recommended to avoid contamination (final concentration 0.05%-0.1%). It is toxic to cells and should be disposed of properly. **Avoid repeated freeze-thaw cycles.**

Background

Tumor necrosis factor receptor superfamily, member 1B (TNFRSF1B), also known as Tumor necrosis factor receptor 2 (TNFR2) or CD120b antigen, is a member of the tumor necrosis factor receptor superfamily. TNFR2/CD120b/TNFRSF1B is a member of the TNF-receptor superfamily. This protein and TNF-receptor 1 form a heterocomplex that mediates the recruitment of two anti-apoptotic proteins, c-IAP1 and c-IAP2, which possess E3 ubiquitin ligase activity. Knockout studies in mice also suggest a role of this protein in protecting neurons from apoptosis by stimulating antioxidative pathways. TNFR2/CD120b/TNFRSF1B is not a major contributing factor to the genetic risk of type 2 diabetes, its associated peripheral neuropathy and hypertension and related metabolic traits in North Indians. Tumor necrosis factor receptor superfamily, member 1B (TNFRSF1B) has been reported to be associated with SLE risk in Japanese populations. TNFR2/CD120b/TNFRSF1B serves as a receptor with high affinity for TNFSF2 and approximately 5-fold lower affinity for homotrimeric TNFSF1. This receptor mediates most of the metabolic effects of TNF-alpha. Isoform 2 blocks TNF-alpha-induced apoptosis, which suggests that it regulates TNF-alpha function by antagonizing its biological activity.

Reference

Komata T, et al., (1999) Association of tumor necrosis factor receptor 2 (TNFR2) polymorphism with susceptibility to systemic lupus erythematosus. *Tissue Antigens.* 53 (6): 527-33.

Tsuchiya N, et al., (2001) Analysis of the association of HLA-DRB1, TNFalpha promoter and TNFR2 (TNFRSF1B) polymorphisms with SLE using transmission disequilibrium test. *Genes Immun.* 2 (6): 317-22.

Guo G, et al., (1999) Role of TNFR1 and TNFR2 receptors in tubulointerstitial fibrosis of obstructive nephropathy. *Am J Physiol.* 277 (5): 766-72.

TNFR2 / CD120b / TNFRSF1B Neutralizing Antibody

Catalog Number: 10417-R00N6




Sino Biological
Biological Solution Specialist

EliteRmab® is a registered trademark of Sino Biological Inc.

Applications

Block – In a functional ELISA which immobilized recombinant Human TNFR2 (Catalog#10417-H08H) at 1 μ g/mL (100 μ L/well) in the plate, the Rabbit anti-Human TNFR2 Monoclonal Antibody (Catalog#10417-R00N6) can block the binding of 2 μ g/mL of biotinylated Human TNF α (Catalog#10602-HNAE) to Human TNFR2, the EC50 is 2.07 μ g/mL.

Neutralization – The neutralization activity of TNFR2/TNFRSF1B Neutralizing Antibody is Measured by its ability to neutralize TNFR2/TNFRSF1B-mediated inhibition of cytotoxicity in the L-929 mouse fibroblast cell line. The Neutralization titer (IC50) is typically 1-3 μ g/mL in the presence of 1 μ g/mL Recombinant Human TNFR2/TNFRSF1B, 0.25 ng/mL Recombinant Human TNF α , and 1 μ g/mL actinomycin D.



TNFR2/TNFRSF1B-mediated inhibition of cytotoxicity was Neutralized by Human TNFR2 Antibody. Recombinant Human TNFR2/TNFRSF1B (Catalog#0417-H08H) inhibits Recombinant Human TNF α (Catalog#10602-HNAE) induced cytotoxicity in the L-929 mouse fibroblast cell line. Inhibition of Recombinant Human TNF α (0.25 ng/mL) activity elicited by Recombinant Human TNFR2/TNFRSF1B (1 μ g/mL) is neutralized by increasing concentrations of Human TNFR2/TNFRSF1B Monoclonal Antibody (Catalog#0417-R00N6). The IC50 is typically 1-3 μ g/mL in the presence of the metabolic inhibitor actinomycin D (1 μ g/mL).