

Cystatin C / CST3 Antibody, Rabbit PAb, Antigen Affinity Purified



Sino Biological
Biological Solution Specialist

Catalog Number: 10439-RP02

GENERAL INFORMATION	
Immunogen:	Recombinant Human Cystatin C / CST3 protein (Catalog#10439-H08H)
Preparation	Produced in rabbits immunized with purified, recombinant Human Cystatin C / CST3 (rh Cystatin C / CST3; Catalog#10439-H08H; NP_000090.1; Ser 27-Ala 146). Cystatin C / CST3 specific IgG was purified by Human Cystatin C / CST3 affinity chromatography.
Ig Type:	Rabbit IgG
Specificity:	Human Cystatin C / CST3
Formulation:	0.2 µm filtered solution in PBS
Storage:	This antibody can be stored at 2°C-8°C for one month without detectable loss of activity. Antibody products are stable for twelve months from date of receipt when stored at -20°C to -80°C. Preservative-Free. Sodium azide is recommended to avoid contamination (final concentration 0.05%-0.1%). It is toxic to cells and should be disposed of properly. Avoid repeated freeze-thaw cycles.
APPLICATIONS	
Applications:	WB,ELISA,IP
RECOMMENDED CONCENTRATION	
Western Blot	WB: 1:500-1:2000
Immunoprecipitation	IP: 2-4 µL/mg of lysate
ELISA	ELISA: 1:5000-1:10000 This antibody can be used at 1:5000-1:10000 with the appropriate secondary reagents to detect Human CST3.


Please Note: Optimal concentrations/dilutions should be determined by the end user.

Cystatin C / CST3 Antibody, Rabbit PAb, Antigen Affinity Purified



Sino Biological
Biological Solution Specialist

Catalog Number: 10439-RP02



Anti-CST3 rabbit polyclonal antibody at 1:500 dilution
Lane A: HepG2 Whole Cell Lysate


Lysates/proteins at 30 µg per lane.

Secondary

Goat Anti-Rabbit IgG H&L (Dylight800) at 1/10000 dilution.

Developed using the Odyssey technique.
Performed under reducing conditions.

Predicted band size: 13 kDa
Observed band size: 13 kDa



CST3 was immunoprecipitated using:
Lane A: 0.5 mg HepG2 Whole Cell Lysate

2 µL anti-CST3 rabbit polyclonal antibody and
15 µL of 50 % Protein G agarose.

Primary antibody:

Anti-CST3 rabbit polyclonal antibody, at 1:200 dilution

Secondary antibody:

Dylight 800-labeled antibody to rabbit IgG (H+L), at 1:5000 dilution

Developed using the odssey technique.
Performed under reducing conditions.

Predicted band size: 14 kDa
Observed band size: 14 kDa