
IL17RA / CD217 Antibody, Mouse MAb







| GENERAL INFORMATION | |
|---------------------------|--|
| Immunogen: | Recombinant Human IL17RA protein (Catalog#10895-H08H) |
| Preparation | This antibody was produced from a hybridoma resulting from the fusion of a mouse myeloma with B cells obtained from a mouse immunized with purified, recombinant Human IL17RA (rh IL17RA; Catalog#10895-H08H; NP_055154.3; Met 1-Trp 320). The IgG fraction of the cell culture supernatant was purified by Protein A affinity chromatography. |
| lg Type: | Mouse IgG1 |
| Clone ID: | 1H3C5 |
| Specificity: | Human IL17RA / CD217 |
| Formulation: | 0.2 µm filtered solution in PBS with 10% Trehalose, pH7.0 |
| Storage: | This antibody can be stored at 2°C-8°C for one month without detectable loss of activity. Antibody products are stable for twelve months from date of receipt when stored at -20°C to -80°C. Preservative-Free. Avoid repeated freeze-thaw cycles. |
| APPLICATIONS | |
| Applications: | ELISA,ELISA(Det),FCM |
| RECOMMENDED CONCENTRATION | |
| Flow Cytometry | FCM: 1:25-1:100 |
| ELISA | ELISA: 1:1000-1:2000 This antibody can be used at 1:1000-1:2000 with the appropriate secondary reagents to detect Human IL17RA. |
| ELISA(Det) | ELISA(Det): 1:1000-1:10000 |

Please Note: Optimal concentrations/dilutions should be determined by the end user.

IL17RA / CD217 Antibody, Mouse MAb







Flow cytometric analysis of Human IL17RA(CD217) expression on human whole blood granulocytes. Cells were stained with purified anti-Human IL17RA(CD217), then a FITC-conjugated second step antibody. The fluorescence histograms were derived from gated events with the forward and side light-scatter characteristics of viable granulocytes.

Flow cytometry was performed on a BD FACSCalibur flow cytometry system. Please refer to www.sinobiological.com/Flow-Cytometry-FACS-Protocols-a-750.html for technical protocols.