

Cyclin A1 / CCNA1 Antibody, Mouse MAb



Sino Biological
Biological Solution Specialist

Catalog Number: 11013-MM06

GENERAL INFORMATION

Immunogen:	Recombinant Human Cyclin A1 / CCNA1 protein (Catalog#11013-H07B)
Preparation	This antibody was produced from a hybridoma resulting from the fusion of a mouse myeloma with B cells obtained from a mouse immunized with purified, recombinant Human Cyclin A1 / CCNA1 (rh Cyclin A1 / CCNA1; Catalog#11013-H07B; NP_003905.1; Met1-Gln465). The IgG fraction of the cell culture supernatant was purified by Protein A affinity chromatography.
Ig Type:	Mouse IgG2b
Clone ID:	06
Specificity:	Human Cyclin A1 / CCNA1
Formulation:	0.2 µm filtered solution in PBS
Storage:	This antibody can be stored at 2°C-8°C for one month without detectable loss of activity. Antibody products are stable for twelve months from date of receipt when stored at -20°C to -80°C. Preservative-Free. Avoid repeated freeze-thaw cycles.
Alternative Names:	CT146

APPLICATIONS

Applications:	FCM, ICC/IF
---------------	-------------

RECOMMENDED CONCENTRATION

ICC/IF	ICC/IF: 1:20-1:100
Flow Cytometry	FCM: 1:25-1:100


Please Note: Optimal concentrations/dilutions should be determined by the end user.

Cyclin A1 / CCNA1 Antibody, Mouse MAb




Sino Biological
Biological Solution Specialist

Catalog Number: 11013-MM06



Immunofluorescence staining of Human CCNA1 in HeLa cells. Cells were fixed with 4% PFA, permeabilized with 1% Triton X-100 in PBS, blocked with 10% serum, and incubated with Mouse anti-Human CCNA1 monoclonal antibody (1:60) at 37°C 1 hour. Then cells were stained with the Alexa Fluor® 488-conjugated Goat Anti-mouse IgG secondary antibody (green) and counterstained with DAPI (blue). Positive staining was localized to cytoplasm and Nucleus.



Flow cytometric analysis of Human CCNA1 expression on HeLa cells. The cells were treated according to manufacturer's manual (BD Pharmingen™ Cat. No. 554714), stained with purified anti-Human CCNA1, then a FITC-conjugated second step antibody. The fluorescence histograms were derived from gated events with the forward and side light-scatter characteristics of intact cells.

Flow cytometry was performed on a BD FACSCalibur flow cytometry system. Please refer to www.sinobiological.com/Flow-Cytometry-FACS-Protocols-a-750.html for technical protocols.