

Rhesus DKK-1 / Dkk1 Protein (256 Asn/Gln, His Tag)

Catalog Number: 11089-K08H



Sino Biological
Biological Solution Specialist

General Information

Gene Name Synonym:

DKK1

Protein Construction:

A DNA sequence encoding the rhesus DKK1 (XP_001098844.1) precursor (Met 1-His 266) (256 Asn / Gln) was expressed, fused with a C-terminal polyhistidine tag.

Source: Rhesus

Expression Host: HEK293 Cells

QC Testing

Purity: > 95 % as determined by SDS-PAGE

Bio Activity:

A DNA sequence encoding the rhesus DKK1 (XP_001098844.1) precursor (Met 2-His 266) (256 Asn / Gln) was expressed, fused with a C-terminal polyhistidine tag.

Endotoxin:

< 1.0 EU per µg of the protein as determined by the LAL method

Stability:

Samples are stable for up to twelve months from date of receipt at -70 °C

Predicted N terminal: Thr 32

Molecular Mass:

The recombinant rhesus DKK1 comprises 246 amino acids with a predicted molecular mass of 27.2 kDa. As a result of glycosylation, the apparent molecular mass of rhesus DKK1 is approximately 38-43 kDa in SDS-PAGE under reducing conditions.

Formulation:

Lyophilized from sterile PBS, pH 7.4

Normally 5 % - 8 % trehalose, mannitol and 0.01% Tween80 are added as protectants before lyophilization. Specific concentrations are included in the hardcopy of COA. Please contact us for any concerns or special requirements.

Usage Guide

Storage:


Store it under sterile conditions at -20°C to -80°C upon receiving. Recommend to aliquot the protein into smaller quantities for optimal storage.

Avoid repeated freeze-thaw cycles.

Reconstitution:

Detailed reconstitution instructions are sent along with the products.

SDS-PAGE:



Protein Description

Dickkopf (DKK) family proteins, consisting of DKK-1, DKK-2, DKK-3 and DKK-4, function as secreted Wnt antagonists by inhibiting Wnt coreceptors LRP5/6. DKK-1, DKK-2, and DKK-4 also bind cell surface Kremen-1 or Kremen-2 and promote the internalization of LRP5/6. Dickkopf related protein 1 (DKK-1) was initially identified as an inducer of head formation in *Xenopus* embryos. DKK-1 protein modulates Wnt signaling pathway during embryonic development. Increased levels of DKK-1 are found in the majority of lung cancers, esophageal squamous cell carcinomas, and hormone-resistant breast cancers, while DKK-1 expression is decreased in malignant melanoma and colorectal cancers.

Manufactured By Sino Biological Inc., FOR RESEARCH USE ONLY. NOT FOR USE IN HUMANS.

For US Customer: Fax: 267-657-0217 • Tel: 215-583-7898

Global Customer: Fax :+86-10-5862-8288 • Tel:+86-400-890-9989 • <http://www.sinobiological.com>