# MERS-CoV Spike/S1 Protein (S1 Subunit, aa 1-725, His Tag)

Catalog Number: 40069-V08B1

# Sino Biological Biological Solution Specialist

### **General Information**

# Gene Name Synonym:

coronavirus s1; coronavirus s2; coronavirus spike; cov spike; ncov RBD; ncov s1; ncov s2; ncov spike; RBD; S; s1; Spike RBD

#### **Protein Construction:**

A DNA sequence encoding the spike protein S1 Subunit MERS-CoV (AFS88936.1)(Met1-Glu725) was fused with a polyhistidine tag at the C-terminus

Source: MERS-CoV

**Expression Host:** Baculovirus-Insect Cells

**QC** Testing

Purity: > 90 % as determined by SDS-PAGE

# **Bio Activity:**

1. Measured by its binding ability in a functional ELISA. Immobilized Spike Protein S1 (aa 1-725) (Cat: 40069-V08B1) at 10  $\mu$ g/ml (100  $\mu$ l/well) can bind biotinylated human DPP4 (Cat: 10688-HNCH).

The EC $_{50}$  of of biotinylated DPP4 (Cat: 10688-HNCH) is 0.52-1.22 µg/ml. 2. Measured by its binding ability in a functional ELISA. Immobilized Spike Protein S1 (aa 1-725) (Cat: 40069-V08B1) at 10 µg/ml (100 µl/well) can bind biotinylated Fc-DPP4 (Cat: 10688-H01H). The EC $_{50}$  of

**Endotoxin:** < 1.0 EU per µg of the protein as determined by the LAL

biotinylated Fc-DPP4 (Cat: 10688-H01H) is 0.02-0.06 μg/ml.

method Stability:

Samples are stable for up to twelve months from date of receipt at -70  $^{\circ}\mathrm{C}$ 

Predicted N terminal: Tyr 18

# **Molecular Mass:**

The recombinant spike protein S1 Subunit MERS-CoV comprises 719 amino acids and has a predicted molecular mass of 79.9 kDa. It migrates as an approximately 94 kDa band in SDS-PAGE under reducing conditions.

#### Formulation:

Lyophilized from sterile 20mM Tris, 500mM NaCl, 10% glycerol, pH 7.4.

Normally 5 % - 8 % trehalose, mannitol and 0.01% Tween80 are added as protectants before lyophilization. Specific concentrations are included in the hardcopy of COA. Please contact us for any concerns or special requirements.

# **Usage Guide**

# Storage:


Store it under sterile conditions at  $-20^{\circ}$ C to  $-80^{\circ}$ C upon receiving. Recommend to aliquot the protein into smaller quantities for optimal storage.

Avoid repeated freeze-thaw cycles.

# Reconstitution:

Detailed reconstitution instructions are sent along with the products.

#### SDS-PAGE:



# **Protein Description**

The spike (S) glycoprotein of coronaviruses contains protrusions that will only bind to certain receptors on the host cell. Known receptors bind S1 are ACE2, angiotensin-converting enzyme 2; DPP4, dipeptidyl peptidase-4; APN, aminopeptidase N; CEACAM, carcinoembryonic antigen-related cell adhesion molecule 1; Sia, sialic acid; O-ac Sia, O-acetylated sialic acid. The spike is essential for both host specificity and viral infectivity. The term 'peplomer' is typically used to refer to a grouping of heterologous proteins on the virus surface that function together. The spike (S) glycoprotein of coronaviruses is known to be essential in the binding of the virus to the host cell at the advent of the infection process. It's been reported that SARS-CoV-2 (COVID-19 coronavirus, 2019-nCoV) can infect the human respiratory epithelial cells through interaction with the human ACE2 receptor.

The spike protein is a large type I transmembrane protein containing two subunits, S1 and S2. S1 mainly contains a receptor binding domain (RBD), which is responsible for recognizing the cell surface receptor. S2 contains basic elements needed for the membrane fusion. The S protein plays key parts in the induction of neutralizing-antibody and T-cell responses, as well as protective immunity.

The main functions for the Spike protein are summarized as: Mediate receptor binding and membrane fusion; Defines the range of the hosts and specificity of the virus; Main component to bind with the neutralizing antibody; Key target for vaccine design; Can be transmitted between different hosts through gene recombination or mutation of the receptor binding domain (RBD), leading to a higher mortality rate.

#### References

1.Shen S, et al. (2007) Expression, glycosylation, and modification of the spike (S) glycoprotein of SARS CoV. Methods Mol Biol. 379: 127-35.

2.Du L, et al. (2009) The spike protein of SARS-CoV--a target for vaccine

3.Xiao X, et al. (2004) The SARS-CoV S glycoprotein. Cell Mol Life Sci. 61 (19-20): 2428-30.

and therapeutic development. Nat Rev Microbiol. 7 (3): 226-36.

Manufactured By Sino Biological Inc., FOR RESEARCH USE ONLY. NOT FOR USE IN HUMANS.

For US Customer: Fax: 267-657-0217 

• Tel: 215-583-7898

Global Customer: Fax :+86-10-5862-8288 ● Tel:+86-400-890-9989 ● http://www.sinobiological.com