

Mouse SRC Kinase / Proto-oncogene c-Src Protein (His & GST Tag)

Catalog Number: 50311-M20B



Sino Biological
Biological Solution Specialist

General Information

Gene Name Synonym:

AW259666; pp60c-src

Protein Construction:

A DNA sequence encoding the mouse SRC (NP_001020566.111) (Met 1-Leu 535) was fused with the N-terminal polyhistidine-tagged GST tag at the N-terminus.

Source: Mouse

Expression Host: Baculovirus-Insect Cells

QC Testing

Purity: > 90 % as determined by SDS-PAGE

Bio Activity:

1. The specific activity was determined to be >50 nmol/min/mg using poly [Glu, Tyr] 4:1 as substrate.

Endotoxin:

< 1.0 EU per µg of the protein as determined by the LAL method

Stability:

Samples are stable for up to twelve months from date of receipt at -70 °C

Predicted N terminal: Met

Molecular Mass:

The recombinant mouse SRC/GST chimera consists of 773 amino acids and has a calculated molecular mass of 87.7 kDa. It migrates as an approximately 80 kDa band in SDS-PAGE under reducing conditions.

Formulation:

Supplied as sterile 20 mM Tris, 300 mM NaCl, 25% Glycerol, 0.5 mM TCEP, pH 8.0.

Specific concentrations are included in the hardcopy of COA. Please contact us for any concerns or special requirements.

Usage Guide

Storage:


Store it under sterile conditions at -20°C to -80°C upon receiving. Recommend to aliquot the protein into smaller quantities for optimal storage.

Avoid repeated freeze-thaw cycles.

Reconstitution:

Detailed reconstitution instructions are sent along with the products.

SDS-PAGE:



Protein Description

Proto-oncogene tyrosine-protein kinase SRC is a hydrophobic protein belonging to the SRC family kinase including nine members that is a family of non-receptor tyrosine kinases. SRC protein may exist in different forms: C-SRC and V-SRC. C-SRC is only activated under certain circumstances where it is required such as growth factor signaling, while V-SRC is a constitutively active as opposed to normal SRC (C-SRC). Thus, V-SRC is an instructive example of an oncogene protein kinase whereas C-SRC is a proto-oncogene protein kinase. Inhibition of SRC with NR2A tyrosine phosphorylation mediated by PSD-95 may contribute to the lithium-induced downregulation of NMDA receptor function and provide neuroprotection against excitotoxicity.

References

1. Juan Ma. et al., 2003, Neuroscience Letters. 348 (3): 185-189.
2. Czernilofsky AP. et al., 1980, Nature. 287: 198-203.
3. Beischlag TV. et al., 2002, Molecular and cellular biology. 22 (12): 4319-33.

Manufactured By Sino Biological Inc., FOR RESEARCH USE ONLY. NOT FOR USE IN HUMANS.

For US Customer: Fax: 267-657-0217 • Tel: 215-583-7898

Global Customer: Fax :+86-10-5862-8288 • Tel:+86-400-890-9989 • <http://www.sinobiological.com>