

Mouse ICOS / AILIM / CD278 Protein (Fc Tag)



Sino Biological
Biological Solution Specialist

Catalog Number: 50466-M02H

General Information

Gene Name Synonym:

AILIM; CCLP; CRP-1; H4; Ly115

Protein Construction:

A DNA sequence encoding the mouse ICOS (Q9WVS0) extracellular domain (Met 1-Leu 142) was fused with the Fc region of human IgG1 at the C-terminus.

Source: Mouse

Expression Host: HEK293 Cells

QC Testing

Purity: > 95 % as determined by SDS-PAGE

Bio Activity:

Immobilized Recombinant Mouse ICOS Ligand / B7-H2 / ICOSLG Protein (His Tag)(Cat:50190-M08H) at 2 µg/ml (100 µl/well) can bind Recombinant Mouse ICOS / AILIM / CD278 Protein (Fc Tag)(Cat:50466-M02H), The EC₅₀ is 100-300 ng/mL.

Endotoxin:

< 1.0 EU per µg of the protein as determined by the LAL method

Predicted N terminal: Glu 21

Molecular Mass:

The recombinant mouse ICOS consists of 363 amino acids and has a predicted molecular mass of 40.9 kDa.

Formulation:

Lyophilized from sterile PBS, pH 7.4

Normally 5 % - 8 % trehalose, mannitol and 0.01% Tween80 are added as protectants before lyophilization. Specific concentrations are included in the hardcopy of COA. Please contact us for any concerns or special requirements.

Usage Guide

Stability & Storage:

Samples are stable for twelve months from date of receipt at -20°C to -80°C.


Store it under sterile conditions at -20°C to -80°C upon receiving. Recommend to aliquot the protein into smaller quantities for optimal storage.

Avoid repeated freeze-thaw cycles.

Reconstitution:

Detailed reconstitution instructions are sent along with the products.

SDS-PAGE:



Protein Description

Inducible costimulator (ICOS), also called AILIM (Activation-Inducible Lymphocyte Immunomediatory Molecule) is a cell-surface receptor and belongs to the CD28 family of immune costimulatory receptors consisting of CD28, CTLA-4, and PD-1. The interaction of B7-H2/ICOS plays a critical role in Th cell differentiation, T-B cell interactions which are essential for the germinal center formation, and humoral immune responses, and as well as the production of cytokine IL-4. Also, ICOS is more potent in the induction of IL-10 production, a cytokine important for the suppressive function of T regulatory cells. The B7-1/B7-2-CD28/CTLA-4 and ICOS-B7RP-1 pathway provide key second signals that can regulate the activation, inhibition, and fine-tuning of T-lymphocyte responses. ICOS stimulates both Th1 and Th2 cytokine production but may have a preferential role in Th2 cell development. Moreover, The B7-1/B7-2-CD28/CTLA-4 and ICOS-B7RP-1 pathway has been suggested as being involved in the development of airway inflammation and airway hyperresponsiveness.

References

- 1.Coyle AJ, *et al.* (2004) The role of ICOS and other costimulatory molecules in allergy and asthma. Springer Semin Immunopathol. 25(3-4): 349-59.
- 2.Chen YQ, *et al.* (2006) CD28/CTLA-4-CD80/CD86 and ICOS-B7RP-1 costimulatory pathway in bronchial asthma. Allergy. 61(1): 15-26.
- 3.van Berkel ME, *et al.* (2006) CD28 and ICOS: similar or separate costimulators of T cells Immunol Lett. 105(2): 115-22.

For Research Use Only. Not for use in diagnostic or therapeutic procedures.

Tel: +86-400-890-9989 (Global), +1-215-583-7898 (USA), +49(0)6196 9678656 (Europe)

Website: <http://www.sinobiological.com>