
HPGD / 15-PGDH Antibody, Rabbit PAb, Antigen Affinity Purified






GENERAL INFORMATION	
Immunogen:	Recombinant Mouse HPGD / 15-PGDH protein (Catalog#50531-M08E)
Preparation	Produced in rabbits immunized with purified, recombinant Mouse HPGD / 15-PGDH (rM HPGD / 15-PGDH; Catalog#50531-M08E; Q8VCC1; Met 1-Ser 269). HPGD / 15-PGDH specific IgG was purified by Mouse HPGD / 15-PGDH affinity chromatography.
Ig Type:	Rabbit IgG
Specificity:	Mouse HPGD / 15-PGDH
Formulation:	0.2 µm filtered solution in PBS
Storage:	This antibody can be stored at $2^{\circ}\text{C-8}^{\circ}\text{C}$ for one month without detectable loss of activity. Antibody products are stable for twelve months from date of receipt when stored at -20°C to -80°C . Preservative-Free. Sodium azide is recommended to avoid contamination (final concentration 0.05%-0.1%). It is toxic to cells and should be disposed of properly. Avoid repeated freeze-thaw cycles.
APPLICATIONS	
Applications:	WB,ELISA,IP
RECOMMENDED CONCENTRATION	
Western Blot	WB: 1:500-1:2000
Immunoprecipitation	IP: 1-4 µL/mg of lysate
ELISA	ELISA: 1:5000-1:10000 This antibody can be used at 1:5000-1:10000 with the appropriate secondary reagents to detect Mouse HPGD.

Please Note: Optimal concentrations/dilutions should be determined by the end user.

HPGD / 15-PGDH Antibody, Rabbit PAb, Antigen Affinity Purified

Catalog Number: 50531-RP02






Anti-HPGD rabbit polyclonal antibody at 1:500 dilution

Lane A: Lovo Whole Cell Lysate

Lysates/proteins at 30 µg per lane. Secondary Goat Anti-Rabbit IgG H&L (Dylight800) at 1/10000 dilution.

Developed using the Odyssey technique. Performed under reducing conditions.

Predicted band size:29 kDa Observed band size:34 kDa (We are unsure as to the identity of these extra bands.)



Mouse HPGD was immunoprecipitated using: Lane A:0.5 mg Caco-2 Whole Cell Lysate

2 μL anti-Mouse HPGD rabbit polyclonal antibody and 15 μl of 50 % Protein G agarose.

Primary antibody: Anti-Mouse HPGD rabbit polyclonal antibody,at 1:200 dilution

Secondary antibody: Dylight 800-labeled antibody to rabbit IgG (H+L), at 1:5000 dilution

Developed using the odssey technique. Performed under reducing conditions.

Predicted band size: 29 kDa Observed band size: 29 kDa