

Mouse CD33 / Siglec-3 Protein (Fc Tag)



Sino Biological
Biological Solution Specialist

Catalog Number: 50712-M02H

General Information

Gene Name Synonym:

gp67; Siglec-3

Protein Construction:

A DNA sequence encoding the extracellular domain of mouse CD33 (Q63994-1) (Met 1-Glu 240) was fused with the Fc region of human IgG1 at the C-terminus.

Source: Mouse

Expression Host: HEK293 Cells

QC Testing

Purity: ≥ 90 % as determined by SDS-PAGE. ≥ 90 % as determined by SEC-HPLC.

Endotoxin:

< 1.0 EU per µg of the protein as determined by the LAL method

Stability:

Samples are stable for up to twelve months from date of receipt at -70 °C

Predicted N terminal: Gln 17

Molecular Mass:

The secreted recombinant mouse CD33/Fc is a disulfide-linked homodimer. The reduced monomer comprises 465 amino acids and has a calculated molecular mass of 51.6 kDa. As a result of glycosylation, the apparent molecular mass of rm CD33/Fc monomer is approximately 55-60 kDa in SDS-PAGE under reducing conditions.

Formulation:

Lyophilized from sterile PBS, pH 7.4

Normally 5 % - 8 % trehalose, mannitol and 0.01% Tween80 are added as protectants before lyophilization. Specific concentrations are included in the hardcopy of COA. Please contact us for any concerns or special requirements.

Usage Guide

Storage:


Store it under sterile conditions at -20°C to -80°C upon receiving. Recommend to aliquot the protein into smaller quantities for optimal storage.

Avoid repeated freeze-thaw cycles.

Reconstitution:

Detailed reconstitution instructions are sent along with the products.

SDS-PAGE:



Protein Description

Myeloid cell surface antigen CD33 also known as Sialic acid binding Ig-like lectin 3, CD33 antigen or Siglec-3, is a member of the immunoglobulin superfamily and SIGLEC (sialic acid binding Ig-like lectin) family. This Single-pass type I membrane protein contains 1 Ig-like C2-type (immunoglobulin-like) domain and 1 Ig-like V-type (immunoglobulin-like) domain. CD33 /Siglec-3 is a putative adhesion molecule of myelomonocytic-derived cells that mediates sialic-acid dependent binding to cells. CD33 /Siglec-3 preferentially binds to alpha-2,6-linked sialic acid. The sialic acid recognition site may be masked by cis interactions with sialic acids on the same cell surface. In the immune response, may act as an inhibitory receptor upon ligand induced tyrosine phosphorylation by recruiting cytoplasmic phosphatase(s) via their SH2 domain(s) that block signal transduction through dephosphorylation of signaling molecules. CD33/Siglec-3 induces apoptosis in acute myeloid leukemia (in vitro). CD33/Siglec-3 can function as a sialic acid-dependent cell adhesion molecule and that binding can be modulated by endogenous sialoglycoconjugates when CD33 is expressed in a plasma membrane.

References

1. Simmons D, *et al.* (1988) Isolation of a cDNA encoding CD33, a differentiation antigen of myeloid progenitor cells. *J Immunol.* 141(8): 2797-800.
2. Ulyanova T, *et al.* (1999) The sialoadhesin CD33 is a myeloid-specific inhibitory receptor. *Eur J Immunol.* 29(11): 3440-9.
3. Freeman SD, *et al.* (1995) Characterization of CD33 as a new member of the sialoadhesin family of cellular interaction molecules. *Blood.* 85(8): 2005-12.

Manufactured By Sino Biological Inc., FOR RESEARCH USE ONLY. NOT FOR USE IN HUMANS.

For US Customer: Fax: 267-657-0217 • Tel: 215-583-7898

Global Customer: Fax :+86-10-5862-8288 • Tel:+86-400-890-9989 • <http://www.sinobiological.com>