

Mouse FLT-3 / CD135 / FLK-2 Protein (His Tag)

Catalog Number: 51119-M08H



Sino Biological
Biological Solution Specialist

General Information

Gene Name Synonym:

B230315G04; CD135; Flk-2; Flk2; Flt-3; Ly72; wml

Protein Construction:

A DNA sequence encoding the mouse FLT3 (EDL05832.1) (Met1-Ser544) was expressed with a C-terminal polyhistidine tag.

Source: Mouse

Expression Host: HEK293 Cells

QC Testing

Purity: ≥ 75 % as determined by SDS-PAGE. ≥ 95 % as determined by SEC-HPLC.

Endotoxin:

< 1.0 EU per µg of the protein as determined by the LAL method

Predicted N terminal: Asn 28

Molecular Mass:

The recombinant mouse FLT3 comprises 528 amino acids and has a predicted molecular mass of 59.5 kDa. As a result of glycosylation, it migrates as an approximately 88 kDa band in SDS-PAGE under reducing conditions.

Formulation:

Lyophilized from sterile PBS, pH 7.4.

Normally 5 % - 8 % trehalose, mannitol and 0.01% Tween80 are added as protectants before lyophilization. Specific concentrations are included in the hardcopy of COA. Please contact us for any concerns or special requirements.

Usage Guide

Stability & Storage:

Samples are stable for twelve months from date of receipt at -20°C to -80°C.

Store it under sterile conditions at -20°C to -80°C upon receiving. Recommend to aliquot the protein into smaller quantities for optimal storage.

Avoid repeated freeze-thaw cycles.

Reconstitution:

Detailed reconstitution instructions are sent along with the products.

SDS-PAGE:



Protein Description

The cluster of differentiation (CD) system is commonly used as cell markers in Immunophenotyping. Different kinds of cells in the immune system can be identified through the surface CD molecules which associating with the immune function of the cell. There are more than 320 CD unique clusters and subclusters have been identified. Some of the CD molecules serve as receptors or ligands important to the cell through initiating a signal cascade which then alter the behavior of the cell. Some CD proteins do not take part in cell signal process but have other functions such as cell adhesion. CD135, also known as FLT-3, FLK-2, is a member of the CD system. CD135 is an important cell surface marker recognized by specific sets of antibodies to identify the types of hematopoietic (blood) progenitors in the bone marrow and it function to differentiate hematopoietic stem cells, which are CD135 negative, from multipotent progenitors, which are CD135 positive. CD135 is a receptor tyrosine kinase type III for the cytokine Flt3 ligand and activate signaling through second messengers by binding to Flt3. Signaling through CD135 is important for lymphocyte development. The encoding gene CD135 is a proto-oncogene to which mutations happened will lead to cancer such as leukemia.

References

1. Zola H, *et al.* (2007) CD molecules 2006-human cell differentiation molecules. *J Immunol Methods*. 318 (1-2): 1-5.
2. Ho IC, *et al.* (2009) GATA3 and the T-cell lineage: essential functions before and after T-helper-2-cell differentiation. *Nat Rev Immunol*. 9 (2): 125-35.
3. Matesanz-Isabel J, *et al.* (2011) New B-cell CD molecules. *Immunology Letters*. 134 (2): 104-12.

For Research Use Only. Not for use in diagnostic or therapeutic procedures.

Tel: +86-400-890-9989 (Global), +1-215-583-7898 (USA), +49(0)6196 9678656 (Europe)

Website: <http://www.sinobiological.com>