

Rat TrkB / NTRK2 Protein (His Tag)

Catalog Number: 80243-R08H



Sino Biological
Biological Solution Specialist

General Information

Gene Name Synonym:

NTRK2

Protein Construction:

A DNA sequence encoding the rat NTRK2 (Q63604-Isoform T1) (Met1-His429) was expressed with a polyhistidine tag at the C-terminus.

Source: Rat

Expression Host: HEK293 Cells

QC Testing

Purity: > 95 % as determined by SDS-PAGE

Bio Activity:

Measured by its binding ability in a functional ELISA. Immobilized rat NTRK2-His at 10 µg/ml (100 µl/well) can bind biotinylated mouse BDNF (cat:50240-MNAS). The EC₅₀ of biotinylated mouse BDNF (cat:50240-MNAS) is 12.2-28.6 ng/ml.

Endotoxin:

< 1.0 EU per µg of the protein as determined by the LAL method

Stability:

Samples are stable for up to twelve months from date of receipt at -70 °C

Predicted N terminal: Cys 32

Molecular Mass:

The recombinant rat NTRK2 comprises 409 amino acids and predicts a molecular mass of 47.1 kDa. The apparent molecular mass of the recombinant protein is approximately 67 kDa in SDS-PAGE under reducing conditions due to glycosylation.

Formulation:

Lyophilized from sterile PBS, pH 7.4

Normally 5 % - 8 % trehalose, mannitol and 0.01% Tween80 are added as protectants before lyophilization. Specific concentrations are included in the hardcopy of COA. Please contact us for any concerns or special requirements.

Usage Guide

Storage:


Store it under sterile conditions at -20°C to -80°C upon receiving. Recommend to aliquot the protein into smaller quantities for optimal storage.

Avoid repeated freeze-thaw cycles.

Reconstitution:

Detailed reconstitution instructions are sent along with the products.

SDS-PAGE:



Protein Description

TrkB receptor also known as TrkB tyrosine kinase or BDNF/NT-3 growth factors receptor or neurotrophic tyrosine kinase, receptor, type 2 (NTRK2) is a single transmembrane catalytic receptors with intracellular tyrosine kinase activity. TrkB/NTRK2 is a member of the neurotrophic tyrosine receptor kinase (NTRK) family. TrkB tyrosine kinase (TrkB) or NTRK2 is coupled to the Ras, Cdc42/Rac/RhoG, MAPK, PI3-K and PLCgamma signaling pathways. There are four members of the Trk family; TrkA, TrkB and TrkC and a related p75NTR receptor. Each family member binds different neurotrophins with varying affinities. TrkB/NTRK has highest affinity for brain-derived neurotrophic factor (BDNF) and is involved in neuronal plasticity, longterm potentiation and apoptosis of CNS neurons. Other neurotrophins include , neurotrophin-3 and neurotrophin-4. TrkB/NTRK is a membrane-bound receptor that, upon neurotrophin binding, phosphorylates itself and members of the MAPK pathway. Signalling through this kinase leads to cell differentiation. Mutations in TrkB/NTRK have been associated with obesity and mood disorders.

References

- 1.Klein R, *et al.* (1990) The trkB tyrosine protein kinase gene codes for a second neurogenic receptor that lacks the catalytic kinase domain. *Cell.* 61 (4): 647-56.
- 2.Rose CR, *et al.* (2003) Truncated TrkB-T1 mediates neurotrophin-evoked calcium signalling in glia cells. *Nature.* 426 (6962): 74-8.
- 3.Yamada K, *et al.* (2004) Brain-derived neurotrophic factor/TrkB signaling in memory processes. *J Pharmacol Sci.* 91 (4): 267-70.

Manufactured By Sino Biological Inc., FOR RESEARCH USE ONLY. NOT FOR USE IN HUMANS.

For US Customer: Fax: 267-657-0217 • Tel: 215-583-7898

Global Customer: Fax :+86-10-5862-8288 • Tel:+86-400-890-9989 • <http://www.sinobiological.com>