

Human IL-12 (IL12A & IL12B Heterodimer) Protein

Catalog Number: CT050-HNAH



Sino Biological
Biological Solution Specialist

General Information

Gene Name Synonym:

CLMF; CLMF2; IL-12A; IL-12B; IL12; IL12 p70; IMD28; IMD29; NFSK; NKSF; NKSF1; NKSF2; P35

Protein Construction:

A DNA sequence encoding the human IL12A (P29459-1)(Met1-Ser219) was expressed, constructed the plasmid 1; A DNA sequence encoding the human IL12B (NP_002178.2)(Met1-Ser328) was expressed, constructed the plasmid 2. The two plasmids were co-expressed and the human IL12A&IL12B heterodimer was purified.

Source: Human

Expression Host: HEK293 Cells

QC Testing

Purity: ≥ 95 % as determined by SDS-PAGE. ≥ 95 % as determined by SEC-HPLC.

Bio Activity:

Measured by its ability to induce Interferon gamma secretion by human natural killer lymphoma NK-92 cells. The ED₅₀ for this effect is 0.2-2 ng/mL.

Endotoxin:

< 5 EU per mg of the protein.

Predicted N terminal: Arg23&Ile23

Molecular Mass:

The recombinant heterodimer of human IL12A&IL12B comprises 503(197+306) amino acids and has a calculated molecular mass of 57.2(22.5+34.7) KDa. The apparent molecular mass of human IL12 heterodimer is approximately 34 & 41 kDa respectively in SDS-PAGE under reducing conditions.

Formulation:

Lyophilized from sterile PBS, pH 7.4.

Normally 5 % - 8 % trehalose, mannitol and 0.01% Tween80 are added as protectants before lyophilization. Specific concentrations are included in the hardcopy of COA. Please contact us for any concerns or special requirements.

Usage Guide

Stability & Storage:

Samples are stable for twelve months from date of receipt at -20°C to -80°C.

Store it under sterile conditions at -20°C to -80°C upon receiving. Recommend to aliquot the protein into smaller quantities for optimal storage.

Avoid repeated freeze-thaw cycles.

Reconstitution:

Detailed reconstitution instructions are sent along with the products.

SDS-PAGE:

