




Mouse Anti-Human IgG₁ Fc

Cat. No.	Format	Size
9054-01	Purified (UNLB)	0.5 mg
9054-04	Alkaline Phosphatase (AP)	1.0 mL
9054-05	Horseradish Peroxidase (HRP)	1.0 mL
9054-08	Biotin (BIOT)	0.5 mg
9054-09	R-phycoerythrin (PE)	0.1 mg



FLISA plate was coated with purified human IgG₁, IgG₂, IgG₃, and IgG₄. Immunoglobulins were detected with serially diluted Mouse Anti-Human IgG₁ Fc-PE (SB Cat. No. 9054-09).

Overview

Clone	HP6001
Isotype	Mouse (BALB/c) IgG _{2b} K
Immunogen	Human IgG ₁ myeloma protein
Specificity	Human IgG ₁ Fc; Mr 146 kDa

Applications

ELISA – Quality tested ²
 FLISA – Quality tested
 FC ⁷
 WB – Reported in literature ³
 Multiplex – Reported in literature ^{4-6,8}

Working Dilutions

ELISA	Purified (UNLB) antibody	≤ 1 µg/mL
	AP conjugate	1:2,000 – 1:8,000
	HRP conjugate	1:4,000 – 1:8,000
	BIOT conjugate	1:5,000 – 1:20,000
FLISA	PE conjugate	≤ 1 µg/mL

Other Applications Since applications vary, you should determine the optimum working dilution for the product that is appropriate for your specific need.

For Research Use Only. Not for Diagnostic or Therapeutic Use.

Handling and Storage

- The purified (UNLB) antibody is supplied as 0.5 mg of purified immunoglobulin in 1.0 mL of borate buffered saline, pH 8.2. *No preservatives or amine-containing buffer salts added.* Store at 2-8°C.
- The alkaline phosphatase (AP) conjugate is supplied as 1.0 mL in a stock solution of 50 mM Tris/1 mM MgCl₂/50% glycerol, pH 8.0, containing NaN₃ as preservative. Store at 2-8°C or long-term at -20°C.
- The horseradish peroxidase (HRP) conjugate is supplied as 1.0 mL in a stock solution of 50% glycerol/50% PBS, pH 7.4. No preservative added. Store at 2-8°C or long-term at -20°C.
- The biotin (BIOT) conjugate is supplied as 0.5 mg in 1.0 mL of PBS/NaN₃. Store at 2-8°C.
- The R-phycoerythrin (PE) conjugate is supplied as 0.1 mg in 0.2 mL of PBS/NaN₃ and a stabilizing agent. Store at 2-8°C. **Do not freeze!**
- Protect fluorochrome-conjugated forms from light. Reagents are stable for the period shown on the label if stored as directed.

Warning

Some reagents contain sodium azide. Please refer to product specific SDS.

References

1. Reimer CB, Phillips DJ, Aloisio CH, Moore DD, Galland GG, Wells TW, et al. Evaluation of thirty-one mouse monoclonal antibodies to human IgG epitopes. *Hybridoma*. 1984;3:263-75. (Immunogen)
2. Ferreira AR, Singh B, Cabrera-Mora M, Magri De Souza AC, Queiroz Marques MT, Porto LC, et al. Evaluation of naturally acquired IgG antibodies to a chimeric and non-chimeric recombinant species of *Plasmodium vivax* reticulocyte binding protein-1: lack of association with HLA-DRB1*/DQB1* in malaria exposed individuals from the Brazilian Amazon. *PLoS One*. 2014;9(8):e105828. (ELISA)
3. he Brazilian Amazon. *PLoS One*. 2014;9(8):e105828. (ELISA)
4. Keen J, Serghides L, Ayi K, Patel SN, Ayisi J, van Eijk A, et al. HIV impairs opsonic phagocytic clearance of pregnancy-associated malaria parasites. *PLoS Med*. 2007;4(5):e181. (WB)
5. Hönger G, Hopfer H, Arnold M, Spriewald BM, Schaub S, Amico P. Pretransplant IgG subclasses of donor-specific human leukocyte antigen antibodies and development of antibody-mediated rejection. *Transplantation*. 2011;92:41-7. (Multiplex)
6. Arnold M, Ntokou I, Doxiadis II, Spriewald BM, Boletis JN, Iniotaki AG. Donor-specific HLA antibodies: evaluating the risk for graft loss in renal transplant recipients with isotype switch from complement fixing IgG1/IgG3 to noncomplement fixing IgG2/IgG4 anti-HLA alloantibodies. *Transpl Int*. 2014;27:253-61. (Multiplex)
7. ies. *Transpl Int*. 2014;27:253-61. (Multiplex)
8. Schaub S, Hönger G, Koller MT, Liwski R, Amico P. Determinants of C1q binding in the single antigen bead assay. *Transplantation*. 2014;98:387-93. (Multiplex)
9. Customer supplied data (FC)
10. Lefaucheur C, Viglietti D, Bentlejewski C, van Huyen JD, Vernerey D, Aubert O, et al. IgG donor-specific anti-human HLA antibody subclasses and kidney allograft antibody-mediated injury. *J Am Soc Nephrol*. 2016;27:293-304. (Multiplex)