


## Anti-KLK6 Antibody



### Description

This gene encodes a member of the kallikrein subfamily of the peptidase S1 family of serine proteases. Growing evidence suggests that many kallikreins are implicated in carcinogenesis and some have potential as novel cancer and other disease biomarkers. The encoded preproprotein is proteolytically processed to generate the mature protease. Expression of this protease is regulated by steroid hormones and may be elevated in multiple human cancers and in serum from psoriasis patients. The encoded protease may participate in the cleavage of amyloid precursor protein and alpha-synuclein, thus implicating this protease in Alzheimer's and Parkinson's disease, respectively. This gene is located in a gene cluster on chromosome 19. Alternative splicing results in multiple transcript variants, at least one of which encodes an isoform that is proteolytically processed.

|                           |  |
|---------------------------|--|
| <b>Model</b>              | STJ113882  |
| <b>Host</b>               | Rabbit   |
| <b>Reactivity</b>         | Human, Mouse, Rat  |
| <b>Applications</b>       | WB   |
| <b>Immunogen</b>          | A synthetic peptide corresponding to a sequence within amino acids 100 to the C-terminus of human KLK6 (NP_002765.1).  |
| <b>Gene ID</b>            | <a href="#">5653</a>   |
| <b>Gene Symbol</b>        | <a href="#">KLK6</a>   |
| <b>Dilution range</b>     | WB 1:500 - 1:2000  |
| <b>Tissue Specificity</b> | In fluids, highest levels found in milk of lactating women followed by cerebrospinal fluid, nipple aspirate fluid and breast cyst fluid, Also found in serum, seminal plasma and some amniotic fluids and breast tumor cytosolic |

extracts, Not detected in urine, At the tissue level, highest concentrations found in glandular tissues such as salivary glands followed by lung, colon, fallopian tube, placenta, breast, pituitary and kidney, Not detected in skin, spleen, bone, thyroid, heart, ureter, liver, muscle, en

|   |  |
|---|--|
| <b>Purification</b>                     | Affinity purification  |
| <b>Note</b>                             | For Research Use Only (RUO).   |
| <b>Protein Name</b>                     | Kallikrein-6   |
| <b>Molecular Weight</b>                 | 26.856 kDa   |
| <b>Clonality</b>                        | Polyclonal   |
| <b>Conjugation</b>                      | Unconjugated   |
| <b>Isotype</b>                          | IgG  |
| <b>Formulation</b>                      | PBS with 0.02% sodium azide, 50% glycerol, pH7.3.  |
| <b>Storage Instruction</b>              | Store at -20C. Avoid freeze / thaw cycles.   |
| <b>Database Links</b>                   | <a href="https://www.ncbi.nlm.nih.gov/condensedbook/condensedbook.cgi?acc=HGNC:6367OMIM:602652">HGNC:6367OMIM:602652</a>   |
| <b>Alternative Names</b>                | Kallikrein-6   |
| <b>Function</b>                         | Serine protease which exhibits a preference for Arg over Lys in the substrate P1 position and for Ser or Pro in the P2 position, Shows activity against amyloid precursor protein, myelin basic protein, gelatin, casein and extracellular matrix proteins such as fibronectin, laminin, vitronectin and collagen, Degrades alpha-synuclein and prevents its polymerization, indicating that it may be involved in the pathogenesis of Parkinson disease and other synucleinopathies, May be involved in regulation of axon outgrowth following spinal cord injury, Tumor cells treated with a neutralizing KLK6 antibody migrate less than control cells, suggesting a role in invasion and metastasis, |
| <b>Cellular Localization</b>            | Secreted, Nucleus, nucleolus, Cytoplasm, Mitochondrion, Microsome,   |
| <b>Post-translational Modifications</b> | Inactivated by autolytic cleavage after Arg-80   |