


## Anti-CHD2 Antibody



### Description

The CHD family of proteins is characterized by the presence of chromo (chromatin organization modifier) domains and SNF2-related helicase/ATPase domains. CHD genes alter gene expression possibly by modification of chromatin structure thus altering access of the transcriptional apparatus to its chromosomal DNA template. Alternatively spliced transcript variants encoding distinct isoforms have been found for this gene.

<b>Model</b>	STJ115439
<b>Host</b>	Rabbit
<b>Reactivity</b>	Human
<b>Applications</b>	WB
<b>Immunogen</b>	A synthetic peptide corresponding to a sequence within amino acids 1700 to the C-terminus of human CHD2 (NP_001262.3).
<b>Gene ID</b>	<a href="#">1106</a>
<b>Gene Symbol</b>	<a href="#">CHD2</a>
<b>Dilution range</b>	WB 1:500 - 1:2000
<b>Purification</b>	Affinity purification
<b>Note</b>	For Research Use Only (RUO).
<b>Protein Name</b>	Chromodomain-helicase-DNA-binding protein 2 CHD-2
<b>Molecular Weight</b>	211.344 kDa
<b>Clonality</b>	Polyclonal

<b>Conjugation</b>	Unconjugated
<b>Isotype</b>	IgG
<b>Formulation</b>	PBS with 0.02% sodium azide, 50% glycerol, pH7.3.
<b>Storage Instruction</b>	Store at -20C. Avoid freeze / thaw cycles.
<b>Database Links</b>	<a href="#">HGNC:1917OMIM:602119</a>
<b>Alternative Names</b>	Chromodomain-helicase-DNA-binding protein 2 CHD-2
<b>Function</b>	DNA-binding helicase that specifically binds to the promoter of target genes, leading to chromatin remodeling, possibly by promoting deposition of histone H3,3, Involved in myogenesis via interaction with MYOD1: binds to myogenic gene regulatory sequences and mediates incorporation of histone H3,3 prior to the onset of myogenic gene expression, promoting their expression ,
<b>Cellular Localization</b>	Nucleus

---

**St John's Laboratory Ltd**

**F** +44 (0)207 681 2580

**T** +44 (0)208 223 3081

**W** <http://www.stjohnslabs.com/>

**E** [info@stjohnslabs.com](mailto:info@stjohnslabs.com)