


Anti-MLST8 Antibody



Model	STJ115560
Host	Rabbit
Reactivity	Human, Mouse, Rat
Applications	IF, WB
Immunogen	Recombinant fusion protein containing a sequence corresponding to amino acids 1-326 of human MLST8 (NP_001186103.1).
Gene ID	64223
Gene Symbol	MLST8
Dilution range	WB 1:500 - 1:2000 IF 1:50 - 1:200
Tissue Specificity	Broadly expressed, with highest levels in skeletal muscle, heart and kidney
Purification	Affinity purification
Note	For Research Use Only (RUO).
Protein Name	Target of rapamycin complex subunit LST8 TORC subunit LST8 G protein beta subunit-like Gable Protein GbetaL Mammalian lethal with SEC13 protein 8 mLST8
Molecular Weight	35.876 kDa
Clonality	Polyclonal
Conjugation	Unconjugated
Isotype	IgG
Formulation	PBS with 0.02% sodium azide, 50% glycerol, pH7.3.

Storage Instruction	Store at -20C. Avoid freeze / thaw cycles.
Database Links	HGNC:24825OMIM:612190Reactome:R-HSA-1257604
Alternative Names	Target of rapamycin complex subunit LST8 TORC subunit LST8 G protein beta subunit-like Gable Protein GbetaL Mammalian lethal with SEC13 protein 8 mLST8
Function	<p>Subunit of both mTORC1 and mTORC2, which regulates cell growth and survival in response to nutrient and hormonal signals, mTORC1 is activated in response to growth factors or amino acids, Growth factor-stimulated mTORC1 activation involves a AKT1-mediated phosphorylation of TSC1-TSC2, which leads to the activation of the RHEB GTPase that potentially activates the protein kinase activity of mTORC1, Amino acid-signaling to mTORC1 requires its relocalization to the lysosomes mediated by the Ragulator complex and the Rag GTPases, Activated mTORC1 up-regulates protein synthesis by phosphorylating key regulators of mRNA translation and ribosome synthesis, mTORC1 phosphorylates EIF4EBP1 and releases it from inhibiting the elongation initiation factor 4E (eIF4E), mTORC1 phosphorylates and activates S6K1 at 'Thr-389', which then promotes protein synthesis by phosphorylating PDCD4 and targeting it for degradation, Within mTORC1, LST8 interacts directly with MTOR and enhances its kinase activity, In nutrient-poor conditions, stabilizes the MTOR-RPTOR interaction and favors RPTOR-mediated inhibition of MTOR activity, mTORC2 is also activated by growth factors, but seems to be nutrient-insensitive, mTORC2 seems to function upstream of Rho GTPases to regulate the actin cytoskeleton, probably by activating one or more Rho-type guanine nucleotide exchange factors, mTORC2 promotes the serum-induced formation of stress-fibers or F-actin, mTORC2 plays a critical role in AKT1 'Ser-473' phosphorylation, which may facilitate the phosphorylation of the activation loop of AKT1 on 'Thr-308' by PDK1 which is a prerequisite for full activation, mTORC2 regulates the phosphorylation of SGK1 at 'Ser-422', mTORC2 also modulates the phosphorylation of PRKCA on 'Ser-657',</p>
Cellular Localization	Cytoplasm