


Anti-MSLN Antibody



Description

This gene encodes a preproprotein that is proteolytically processed to generate two protein products, megakaryocyte potentiating factor and mesothelin. Megakaryocyte potentiating factor functions as a cytokine that can stimulate colony formation of bone marrow megakaryocytes. Mesothelin is a glycosylphosphatidylinositol-anchored cell-surface protein that may function as a cell adhesion protein. This protein is overexpressed

Mesothelin is a glycosylphosphatidylinositol-anchored cell-surface protein that may function as a cell adhesion protein. This protein is overexpressed in epithelial mesotheliomas, ovarian cancers and in specific squamous cell carcinomas. Alternative splicing results in multiple transcript variants, at least one of which encodes an isoform that is proteolytically processed.

Model STJ116034

Host Rabbit

Reactivity Human, Rat

Applications IF, IHC, WB

Immunogen Recombinant fusion protein containing a sequence corresponding to amino

acids 37-296 of human MSLN (NP_001170826.1).

Gene ID 10232

Gene Symbol MSLN

Dilution range WB 1:500 - 1:2000

IHC 1:50 - 1:200 IF 1:50 - 1:100

Tissue Specificity Expressed in lung, Expressed at low levels in heart, placenta and kidney,

Expressed in mesothelial cells, Highly expressed in mesotheliomas, ovarian

cancers, and some squamous cell carcinomas (at protein level)

Purification Affinity purification

Note For Research Use Only (RUO).

Protein Name Mesothelin CAK1 antigen Pre-pro-megakaryocyte-potentiating factor

Molecular Weight 68.986 kDa

Clonality Polyclonal

Conjugation Unconjugated

Isotype IgG

Formulation PBS with 0.02% sodium azide, 50% glycerol, pH7.3.

Storage Instruction Store at -20C. Avoid freeze / thaw cycles.

Database Links HGNC:73710MIM:601051Reactome:R-HSA-163125

Alternative Names Mesothelin CAK1 antigen Pre-pro-megakaryocyte-potentiating factor

Function Membrane-anchored forms may play a role in cellular adhesion

Cellular Localization Cell membrane

Post-translational Both MPF and the cleaved form of mesothelin are N-glycosylated,

Modifications

St John's Laboratory Ltd

F +44 (0)207 681 2580 **T** +44 (0)208 223 3081

W http://www.stjohnslabs.com/ E info@stjohnslabs.com