


Anti-TNFRSF19 Antibody



Description

The protein encoded by this gene is a member of the TNF-receptor superfamily. This receptor is highly expressed during embryonic development. It has been shown to interact with TRAF family members, and to activate JNK signaling pathway when overexpressed in cells. This receptor is capable of inducing apoptosis by a caspase-independent mechanism, and it is thought to play an essential role in embryonic development. Alternatively spliced transcript variants encoding distinct isoforms have been described.

Model	STJ116496
Host	Rabbit
Reactivity	Human
Applications	WB
Immunogen	Recombinant fusion protein containing a sequence corresponding to amino acids 30-170 of human TNFRSF19 (NP_061117.2).
Gene ID	55504
Gene Symbol	TNFRSF19
Dilution range	WB 1:500 - 1:1000
Tissue Specificity	Highly expressed in prostate, Detected at lower levels in thymus, spleen, testis, uterus, small intestine, colon and peripheral blood leukocytes
Purification	Affinity purification
Note	For Research Use Only (RUO).
Protein Name	Tumor necrosis factor receptor superfamily member 19 TRADE Toxicity and

	JNK inducer
Molecular Weight	46.015 kDa
Clonality	Polyclonal
Conjugation	Unconjugated
Isotype	IgG
Formulation	PBS with 0.02% sodium azide, 50% glycerol, pH7.3.
Storage Instruction	Store at -20C. Avoid freeze / thaw cycles.
Database Links	HGNC:11915OMIM:606122
Alternative Names	Tumor necrosis factor receptor superfamily member 19 TRADE Toxicity and JNK inducer
Function	Can mediate activation of JNK and NF-kappa-B, May promote caspase-independent cell death
Cellular Localization	Membrane

St John's Laboratory Ltd

F +44 (0)207 681 2580

T +44 (0)208 223 3081

W <http://www.stjohnslabs.com/>

E info@stjohnslabs.com