

## Anti-NCR1 Antibody

---



<b>Model</b>	STJ116755
<b>Host</b>	Rabbit
<b>Reactivity</b>	Human, Mouse
<b>Applications</b>	WB
<b>Immunogen</b>	Recombinant fusion protein containing a sequence corresponding to amino acids 151-303 of human NCR1 (NP_001138929.2).
<b>Gene ID</b>	<a href="#">9437</a>
<b>Gene Symbol</b>	<a href="#">NCR1</a>
<b>Dilution range</b>	WB 1:500 - 1:2000
<b>Tissue Specificity</b>	Selectively expressed by both resting and activated NK cells
<b>Purification</b>	Affinity purification
<b>Note</b>	For Research Use Only (RUO).
<b>Protein Name</b>	Natural cytotoxicity triggering receptor 1 Lymphocyte antigen 94 homolog NK cell-activating receptor Natural killer cell p46-related protein NK-p46 NKp46 hNKp46 CD antigen CD335
<b>Molecular Weight</b>	34.481 kDa
<b>Clonality</b>	Polyclonal
<b>Conjugation</b>	Unconjugated
<b>Isotype</b>	IgG
<b>Formulation</b>	PBS with 0.02% sodium azide, 50% glycerol, pH7.3.
<b>Storage Instruction</b>	Store at -20C. Avoid freeze / thaw cycles.

<b>Database Links</b>	<a href="#">HGNC:6731</a> <a href="#">OMIM:604530</a> <a href="#">Reactome:R-HSA-198933</a>
<b>Alternative Names</b>	Natural cytotoxicity triggering receptor 1 Lymphocyte antigen 94 homolog NK cell-activating receptor Natural killer cell p46-related protein NK-p46 NKp46 hNKp46 CD antigen CD335
<b>Function</b>	Cytotoxicity-activating receptor that may contribute to the increased efficiency of activated natural killer (NK) cells to mediate tumor cell lysis,
<b>Cellular Localization</b>	Cell membrane
<b>Post-translational Modifications</b>	N-glycosylated,

---

**St John's Laboratory Ltd**

**F** +44 (0)207 681 2580

**T** +44 (0)208 223 3081

**W** <http://www.stjohnslabs.com/>

**E** [info@stjohnslabs.com](mailto:info@stjohnslabs.com)