


Anti-myc antibody



Description

Goat polyclonal antibody to myc epitope tag. c-Myc epitope tag is useful for the labelling and detection of proteins by immunoprecipitation, immunostaining and immunoblotting techniques. Because of its small size, it is unlikely to affect the tagged protein's biochemical properties.

Model	STJ140011
Host	Goat
Applications	WB
Immunogen	Purified recombinant peptide EQKLISEEDL corresponding to amino acids 410-419 of Human c-Myc produced in E. coli.
Immunogen Region	EQKLISEEDL, at 410-419 aa
Dilution range	Western blot 1:500-1:2,000 Immunofluorescence ND Immunohistochemistry (paraffin) ND Immunohistochemistry (frozen) ND
Specificity	This antibody recognizes recombinant proteins containing EQKLISEEDL epitope tag fused to either amino- or carboxy-terminal of targeted proteins in transfected cells.
Purification	This antibody is epitope-affinity purified from goat antiserum.
Note	For research use only (RUO).
Protein Name	myc
Clonality	Polyclonal
Conjugation	Unconjugated
Isotype	IgG

Formulation	PBS, 20% glycerol and 0.05% sodium azide.
Concentration	3 mg/mL
Storage Instruction	Store at -20°C, and avoid repeated freeze-thaw cycles.
Alternative Names	myc

St John's Laboratory Ltd

F +44 (0)207 681 2580

T +44 (0)208 223 3081

W <http://www.stjohnslabs.com/>

E info@stjohnslabs.com