


## **Anti-CD19 antibody**



**Description** Goat polyclonal antibody to CD19. CD19 is a type I transmembrane

glycoprotein with 2 C-type Ig domains and multiple phosphorylation sites on long cytoplasmic domain. It is a cell surface molecule which assembles with the antigen receptor of B lymphocytes in order to decrease the threshold for antigen receptor- dependent stimulation. It is a major B

lineage marker.

Model STJ140024

**Host** Goat

**Reactivity** Avian, Bovine, Canine, Donkey, Feline, Goat, Guinea Pig, Hamster, Horse,

Human, Mouse, Other, Porcine, Rabbit, Rat, Sheep, Simian

**Applications** WB

**Immunogen** Purified recombinant peptide within residues 510 aa to the C-terminus of

human CD19 produced in E. coli.

Immunogen Region C-Term

Gene ID 930
Gene Symbol CD19

**Dilution range** Western blot 1:500-1:2,000 Immunofluorescence ND Immunohistochemistry

(paraffin) ND Immunohistochemistry (frozen) ND

**Purification** This antibody is epitope-affinity purified from goat antiserum.

**Note** For research use only (RUO).

Protein Name B-lymphocyte antigen CD19 B-lymphocyte surface antigen B4 Differentiation

antigen CD19 T-cell surface antigen Leu-12 CD antigen CD19

Molecular Weight 62 kDa

Polyclonal **Clonality** 

Unconjugated Conjugation

**IgG Isotype** 

**Formulation** PBS, 20% glycerol and 0.05% sodium azide.

Concentration 3 mg/mL

**Storage Instruction** Store at -20°, and avoid repeated freeze-thaw cycles.

**Database Links** HGNC:1633OMIM:107265

B-lymphocyte antigen CD19 B-lymphocyte surface antigen B4 Differentiation **Alternative Names** 

antigen CD19 T-cell surface antigen Leu-12 CD antigen CD19

**Function** Assembles with the antigen receptor of B-lymphocytes in order to decrease

the threshold for antigen receptor-dependent stimulation.

Cellular Localization Membrane. Single-pass type I membrane protein.

Post-translational Phosphorylated on serine and threonine upon DNA damage, probably by **Modifications** 

ATM or ATR. Phosphorylated on tyrosine following B-cell activation.

Phosphorylated on tyrosine residues by LYN.

{ECO:0000269|PubMed:10706702, ECO:0000269|PubMed:7687428,

ECO:0000269|PubMed:7687539}.

St John's Laboratory Ltd

**F** +44 (0)207 681 2580

T+44 (0)208 223 3081

W http://www.stjohnslabs.com/ E info@stjohnslabs.com