
## **Anti-GroEL antibody**





**Description** Goat polyclonal antibody to GroEL. GroEL protein is a 60 kDa chaperone

that promotes refolding of misfolded polypeptides especially under

stressful conditions.

Model STJ140039

**Host** Goat

**Reactivity** Burkholderia, Other, Pseudomonas

**Applications** WB

**Immunogen** Purified recombinant peptide derived from within residues 400 aa to the C-

terminus of Burkholderia GroEL produced in E. coli.

**Immunogen Region** C-Term

Gene Symbol HSPD1

**Dilution range** Western blot 1:500-1:5,000 Immunofluorescence ND Immunohistochemistry

(paraffin) ND Immunohistochemistry (frozen) ND

**Specificity** The Ab recognizes the endogenous protein in Burkholderia and Pseudomonas

by Western blot.

**Purification** This antibody is epitope-affinity purified from goat antiserum.

**Note** For research use only (RUO).

**Protein Name** 60 kDa heat shock protein, mitochondrial 60 kDa chaperonin Chaperonin 60

CPN60 Heat shock protein 60 HSP-60 Hsp60 HuCHA60 Mitochondrial

matrix protein P1 P60 lymphocyte protein

Molecular Weight 1 kDa

**Clonality** Polyclonal

**Conjugation** Unconjugated

**Isotype** IgG

**Formulation** PBS, 20% glycerol and 0.05% sodium azide.

**Concentration** 2 mg/mL

**Storage Instruction** Store at -20°, and avoid repeated freeze-thaw cycles.

Alternative Names 60 kDa heat shock protein, mitochondrial 60 kDa chaperonin Chaperonin 60

CPN60 Heat shock protein 60 HSP-60 Hsp60 HuCHA60 Mitochondrial

matrix protein P1 P60 lymphocyte protein

**Function** Chaperonin implicated in mitochondrial protein import and macromolecular

assembly. Together with Hsp10, facilitates the correct folding of imported proteins. May also prevent misfolding and promote the refolding and proper assembly of unfolded polypeptides generated under stress conditions in the mitochondrial matrix (PubMed:1346131, PubMed:11422376). The functional units of these chaperonins consist of heptameric rings of the large subunit Hsp60, which function as a back-to-back double ring. In a cyclic reaction, Hsp60 ring complexes bind one unfolded substrate protein per ring, followed by the binding of ATP and association with 2 heptameric rings of the cochaperonin Hsp10. This leads to sequestration of the substrate protein in the inner cavity of Hsp60 where, for a certain period of time, it can fold undisturbed by other cell components. Synchronous hydrolysis of ATP in all

Hsp60 subunits results in the dissociation of the chaperonin rings and the

release of ADP and the folded substrate protein (Probable)

Cellular Localization Mithocondria

St John's Laboratory Ltd

**F** +44 (0)207 681 2580

**T** +44 (0)208 223 3081

W http://www.stjohnslabs.com/ E info@stjohnslabs.com