
Anti-Rab8b antibody



Description

Goat polyclonal antibody to mouse Rab8. Rab8 belongs to the small GTPase superfamily, Rab family. It has cell-type specific function during the process of exocytosis and coordinates regulation of the cytoskeleton. Rab8 also appears to interface with endocytic recycling circuits. At the molecular level, this GTPase regulates both the export of vesicles from the trans- Golgi apparatus, and the directed translocation along actin filaments and microtubules.

Model STJ140088

Host Goat

Reactivity Avian, Bovine, Canine, Donkey, Feline, Goat, Guinea Pig, Hamster, Horse,

Human, Mouse, Other, Porcine, Rabbit, Rat, Sheep, Simian

Applications WB

Immunogen Purified recombinant peptide derived from within residues 107 aa to the C-

terminus of mouse Rab8b produced in E. coli.

Immunogen Region C-Term

Gene ID <u>51762</u>

Gene Symbol RAB8B

Dilution range Western blot 1:250-1:1,000 Immunofluorescence ND Immunohistochemistry

(paraffin) ND Immunohistochemistry (frozen) ND

Purification This antibody is epitope-affinity purified from goat antiserum.

Note For research use only (RUO).

Protein Name Ras-related protein Rab-8B

Molecular Weight 24 kDa

Clonality Polyclonal

Conjugation Unconjugated

Isotype IgG

Formulation PBS, 20% glycerol and 0.05% sodium azide.

Concentration 4 mg/mL

Storage Instruction Store at -20°, and avoid repeated freeze-thaw cycles.

Database Links HGNC:30273OMIM:613532

Alternative Names Ras-related protein Rab-8B

Function The small GTPases Rab are key regulators of intracellular membrane

trafficking, from the formation of transport vesicles to their fusion with membranes. Rabs cycle between an inactive GDP-bound form and an active GTP-bound form that is able to recruit to membranes different sets of downstream effectors directly responsible for vesicle formation, movement, tethering and fusion. That Rab may be involved in polarized vesicular trafficking and neurotransmitter release. May participate in cell junction

dynamics in Sertoli cells.

Cellular Localization Cell membrane Cytoplasmic vesicle, phagosome Cytoplasmic vesicle,

phagosome membrane. Recruited to phagosomes containing S.aureus or

M.tuberculosis.

St John's Laboratory Ltd

F +44 (0)207 681 2580

T +44 (0)208 223 3081

W http://www.stjohnslabs.com/ E info@stjohnslabs.com