


Anti-Rab5a antibody



Description

Goat polyclonal antibody to mouse Rab5a. Rab5a belongs to the small GTPase superfamily, Rab family. Rab5a is an Early Endosome Marker and functions as a key regulator of vesicular trafficking during early endocytosis.

Model STJ140104

Host Goat

Reactivity Avian, Bovine, Canine, Donkey, Feline, Goat, Guinea Pig, Hamster, Horse, Human, Mouse, Other, Porcine, Rabbit, Rat, Sheep, Simian

Applications WB

Immunogen Purified recombinant peptide derived from within residues 115 aa to the C-terminus of Rab5a produced in E. coli.

Immunogen Region C-Term

Gene ID [5868](#)

Gene Symbol [RAB5A](#)

Dilution range Western blot 1:250-1:2,000 Immunofluorescence ND Immunohistochemistry (paraffin) ND Immunohistochemistry (frozen) ND

Specificity Detects Rab5a protein by Western blot in cell lysates and transfected cells with GFP-Rab5a. This Ab is specific for Rab5a; it does not recognize Rab5b and Rab5c.

Purification This antibody is epitope-affinity purified from goat antiserum.

Note For research use only (RUO).

Protein Name Ras-related protein Rab-5A

Molecular Weight	24 kDa
Clonality	Polyclonal
Conjugation	Unconjugated
Isotype	IgG
Formulation	PBS, 20% glycerol and 0.05% sodium azide.
Concentration	3 mg/mL
Storage Instruction	Store at -20°, and avoid repeated freeze-thaw cycles.
Database Links	HGNC:97830 MIM:179512
Alternative Names	Ras-related protein Rab-5A
Function	<p>The small GTPases Rab are key regulators of intracellular membrane trafficking, from the formation of transport vesicles to their fusion with membranes. Rabs cycle between an inactive GDP-bound form and an active GTP-bound form that is able to recruit to membranes different sets of downstream effectors directly responsible for vesicle formation, movement, tethering and fusion. RAB5A is required for the fusion of plasma membranes and early endosomes . Contributes to the regulation of filopodia extension . Required for the exosomal release of SDCBP, CD63, PDZD6IP and syndecan . Regulates maturation of apoptotic cell-containing phagosomes, probably downstream of DYN2 and PIK3C3 .</p>
Cellular Localization	<p>Cell membrane Early endosome membrane Melanosome Cytoplasmic vesicle Cell projection, ruffle Membrane Cytoplasm, cytosol. Cytoplasmic vesicle, phagosome membrane Endosome membrane. Enriched in stage I melanosomes . Alternates between membrane-bound and cytosolic forms (Probable).</p>