
## Anti-Rab9a antibody



**Description** Goat polyclonal antibody to Rab9a. Rab9a belongs to the small GTPase

superfamily, Rab family. This protein may be involved in endosome-to-

Golgi transport.

Model STJ140136

**Host** Goat

**Reactivity** Avian, Bovine, Canine, Donkey, Feline, Goat, Guinea Pig, Hamster, Horse,

Human, Mouse, Other, Porcine, Rabbit, Rat, Sheep, Simian

**Applications** WB

**Immunogen** Purified recombinant peptide derived from within residues 115 aa to the C-

terminus of Rab9a produced in E. coli.

**Immunogen Region** C-Term

**Gene ID** <u>9367</u>

Gene Symbol RAB9A

**Dilution range** Western blot 1:250-1:2,000 Immunofluorescence ND Immunohistochemistry

(paraffin) ND Immunohistochemistry (frozen) ND

**Purification** This antibody is epitope-affinity purified from goat antiserum.

**Note** For research use only (RUO).

**Protein Name** Ras-related protein Rab-9A

**Molecular Weight** 23 kDa

**Clonality** Polyclonal

Conjugation Unconjugated

IgG **Isotype** 

**Formulation** PBS, 20% glycerol and 0.05% sodium azide.

4 mg/mL Concentration

Store at -20°, and avoid repeated freeze-thaw cycles. **Storage Instruction** 

**Database Links** HGNC:9792OMIM:300284

**Alternative Names** Ras-related protein Rab-9A

**Function** Involved in the transport of proteins between the endosomes and the trans

> Golgi network. Involved in the recruitment of SGSM2 to melanosomes and is required for the proper trafficking of melanogenic enzymes TYR, TYRP1 and

DCT/TYRP2 to melanosomes in melanocytes.

Cell membrane Endoplasmic reticulum membrane Golgi apparatus membrane **Cellular Localization** 

> Late endosome Cytoplasmic vesicle, phagosome membrane Cytoplasmic vesicle, phagosome Cytoplasmic vesicle membrane Melanosome. Colocalizes with OSBPL1A at the late endosome. Recruited to phagosomes containing

S.aureus or M.tuberculosis.

St John's Laboratory Ltd

**F** +44 (0)207 681 2580 T+44 (0)208 223 3081

W http://www.stjohnslabs.com/

E info@stjohnslabs.com