

Anti-T2AG antibody



Description	Unconjugated Rabbit polyclonal to T2AG
Model	STJ190199
Host	Rabbit
Reactivity	Human, Mouse, Rat
Applications	ELISA, WB
Immunogen	Synthesized peptide derived from human T2AG protein.
Immunogen Region	30-110aa
Gene ID	2958
Gene Symbol	GTF2A2
Dilution range	WB 1:500-2000 ELISA 1:5000-20000
Specificity	T2AG Polyclonal Antibody detects endogenous levels of protein.
Purification	T2AG antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
Note	For Research Use Only (RUO).
Protein Name	Transcription initiation factor IIA subunit 2 General transcription factor IIA subunit 2 TFIIA p12 subunit TFIIA-12 TFIIAS Transcription initiation factor IIA gamma chain TFIIA-gamma
Molecular Weight	11 kDa
Clonality	Polyclonal
Conjugation	Unconjugated

Isotype	IgG
Formulation	Liquid form in PBS containing 50% glycerol, and 0.02% sodium azide.
Concentration	1 mg/ml
Storage Instruction	Store at -20°C, and avoid repeat freeze-thaw cycles.
Database Links	HGNC:4647 OMIM:600519
Alternative Names	Transcription initiation factor IIA subunit 2 General transcription factor IIA subunit 2 TFIIA p12 subunit TFIIA-12 TFIIAS Transcription initiation factor IIA gamma chain TFIIA-gamma
Function	TFIIA is a component of the transcription machinery of RNA polymerase II and plays an important role in transcriptional activation. TFIIA in a complex with TBP mediates transcriptional activity.
Cellular Localization	Nucleus.

St John's Laboratory Ltd

F +44 (0)207 681 2580

T +44 (0)208 223 3081

W <http://www.stjohnslabs.com/>

E info@stjohnslabs.com