

Anti-TAF10 antibody



Description	Unconjugated Rabbit polyclonal to TAF10
Model	STJ190211
Host	Rabbit
Reactivity	Human, Mouse
Applications	ELISA, WB
Immunogen	Synthesized peptide derived from human TAF10 protein.
Immunogen Region	130-210aa
Gene ID	6881
Gene Symbol	TAF10
Dilution range	WB 1:500-2000 ELISA 1:5000-20000
Specificity	TAF10 Polyclonal Antibody detects endogenous levels of protein.
Purification	TAF10 antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
Note	For Research Use Only (RUO).
Protein Name	Transcription initiation factor TFIID subunit 10 STAF28 Transcription initiation factor TFIID 30 kDa subunit TAF II30 TAFII-30 TAFII30
Molecular Weight	23 kDa
Clonality	Polyclonal
Conjugation	Unconjugated

Isotype	IgG
Formulation	Liquid form in PBS containing 50% glycerol, and 0.02% sodium azide.
Concentration	1 mg/ml
Storage Instruction	Store at -20°C, and avoid repeat freeze-thaw cycles.
Database Links	HGNC:11543 OMIM:600475
Alternative Names	Transcription initiation factor TFIID subunit 10 STAF28 Transcription initiation factor TFIID 30 kDa subunit TAF II30 TAFII-30 TAFII30
Function	TAFs are components of the transcription factor IID (TFIID) complex, PCAF histone acetylase complex and TBP-free TAFII complex (TFTC). TFIID is a multimeric protein complex that plays a central role in mediating promoter responses to various activators and repressors.
Sequence and Domain Family	The [KR]-[STA]-K motif is specifically recognized by the SETD7 methyltransferase.
Cellular Localization	Nucleus
Post-translational Modifications	Monomethylated at Lys-189 by SETD7, leading to increase its affinity for RNA polymerase II.

St John's Laboratory Ltd

F +44 (0)207 681 2580

T +44 (0)208 223 3081

W <http://www.stjohnslabs.com/>

E info@stjohnslabs.com