

Anti-CRLF3 antibody



| | |
|--------------------|-----------------------------------------|
| Description | Unconjugated Rabbit polyclonal to CRLF3 |
|--------------------|-----------------------------------------|

| | |
|---------------------------|-------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Model | STJ190292 |
| Host | Rabbit |
| Reactivity | Human, Mouse |
| Applications | ELISA, WB |
| Immunogen | Synthesized peptide derived from human CRLF3 protein. |
| Immunogen Region | 270-350aa |
| Gene ID | 51379 |
| Gene Symbol | CRLF3 |
| Dilution range | WB 1:500-2000 ELISA 1:5000-20000 |
| Specificity | CRLF3 Polyclonal Antibody detects endogenous levels of protein. |
| Tissue Specificity | Expressed in several embryonic and adult tissues, including adult and fetal brain, liver, spleen and pancreas. Expressed in adult, but not fetal kidney. Expressed in skin and squamous cell carcinoma (SCC) and in several other cancer types. Also detected in lesion actinic keratosis (AK). |
| Purification | CRLF3 antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen. |
| Note | For Research Use Only (RUO). |
| Protein Name | Cytokine receptor-like factor 3 Cytokine receptor-like molecule 9 CREME-9 Cytokine receptor-related protein 4 Type I cytokine receptor-like factor p48 |

| | |
|------------------------------|-----------------------------------------------------------------------------------------------------------------------------------------------------------|
| Molecular Weight | 48 kDa |
| Clonality | Polyclonal |
| Conjugation | Unconjugated |
| Isotype | IgG |
| Formulation | Liquid form in PBS containing 50% glycerol, and 0.02% sodium azide. |
| Concentration | 1 mg/ml |
| Storage Instruction | Store at -20°C, and avoid repeat freeze-thaw cycles. |
| Database Links | HGNC:17177 OMIM:614853 |
| Alternative Names | Cytokine receptor-like factor 3 Cytokine receptor-like molecule 9 CREME-9 Cytokine receptor-related protein 4 Type I cytokine receptor-like factor p48 |
| Function | May play a role in the negative regulation of cell cycle progression. |
| Cellular Localization | Cytoplasm. The EGFP- and MYC-tagged protein has been shown to be cytoplasmic. |

St John's Laboratory Ltd

F +44 (0)207 681 2580

T +44 (0)208 223 3081

W <http://www.stjohnslabs.com/>

E info@stjohnslabs.com