

## Anti-LRP1B antibody

---



|                           |  |
|---------------------------|--|
| <b>Description</b>        | Unconjugated Rabbit polyclonal to LRP1B  |
| <b>Model</b>              | STJ190935  |
| <b>Host</b>               | Rabbit   |
| <b>Reactivity</b>         | Human, Mouse   |
| <b>Applications</b>       | IHC  |
| <b>Immunogen</b>          | Synthesized peptide derived from human LRP1B protein.  |
| <b>Immunogen Region</b>   | 200-280aa  |
| <b>Gene ID</b>            | <a href="#">53353</a>  |
| <b>Gene Symbol</b>        | <a href="#">LRP1B</a>  |
| <b>Dilution range</b>     | IHC-p 1:50-300   |
| <b>Specificity</b>        | LRP1B Polyclonal Antibody detects endogenous levels of protein.  |
| <b>Tissue Specificity</b> | Expressed in thyroid gland and in salivary gland, as well as in adult and fetal brain.   |
| <b>Purification</b>       | LRP1B antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.              |
| <b>Note</b>               | For Research Use Only (RUO).   |
| <b>Protein Name</b>       | Low-density lipoprotein receptor-related protein 1B LRP-1B Low-density lipoprotein receptor-related protein-deleted in tumor LRP-DIT |
| <b>Molecular Weight</b>   | 505 kDa  |
| <b>Clonality</b>          | Polyclonal   |

|                              |  |
|------------------------------|--|
| <b>Conjugation</b>           | Unconjugated   |
| <b>Isotype</b>               | IgG  |
| <b>Formulation</b>           | Liquid form in PBS containing 50% glycerol, and 0.02% sodium azide.  |
| <b>Concentration</b>         | 1 mg/ml  |
| <b>Storage Instruction</b>   | Store at -20°C, and avoid repeat freeze-thaw cycles.   |
| <b>Database Links</b>        | <a href="#">HGNC:6693OMIM:608766</a>   |
| <b>Alternative Names</b>     | Low-density lipoprotein receptor-related protein 1B LRP-1B Low-density lipoprotein receptor-related protein-deleted in tumor LRP-DIT |
| <b>Function</b>              | Potential cell surface proteins that bind and internalize ligands in the process of receptor-mediated endocytosis.                   |
| <b>Cellular Localization</b> | Membrane   |

---

**St John's Laboratory Ltd**

**F** +44 (0)207 681 2580

**T** +44 (0)208 223 3081

**W** <http://www.stjohnslabs.com/>

**E** [info@stjohnslabs.com](mailto:info@stjohnslabs.com)