

Anti-NOG2 antibody



Description	Unconjugated Rabbit polyclonal to NOG2
Model	STJ191110
Host	Rabbit
Reactivity	Human, Mouse
Applications	ELISA, WB
Immunogen	Synthesized peptide derived from human NOG2 protein.
Immunogen Region	670-750aa
Gene ID	29889
Gene Symbol	GNL2
Dilution range	WB 1:500-2000 ELISA 1:5000-20000
Specificity	NOG2 Polyclonal Antibody detects endogenous levels of protein.
Tissue Specificity	Ubiquitously expressed at relatively low levels in all human tissues tested, with the highest level of expression in the testes.
Purification	NOG2 antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
Note	For Research Use Only (RUO).
Protein Name	Nucleolar GTP-binding protein 2 Autoantigen NGP-1
Molecular Weight	80 kDa
Clonality	Polyclonal

Conjugation	Unconjugated
Isotype	IgG
Formulation	Liquid form in PBS containing 50% glycerol, and 0.02% sodium azide.
Concentration	1 mg/ml
Storage Instruction	Store at -20°C, and avoid repeat freeze-thaw cycles.
Database Links	HGNC:29925OMIM:609365
Alternative Names	Nucleolar GTP-binding protein 2 Autoantigen NGP-1
Function	GTPase that associates with pre-60S ribosomal subunits in the nucleolus and is required for their nuclear export and maturation.
Cellular Localization	Nucleus, nucleolus

St John's Laboratory Ltd

F +44 (0)207 681 2580
T +44 (0)208 223 3081

W <http://www.stjohnslabs.com/>
E info@stjohnslabs.com