

## Anti-PRD10 antibody

---



|                         |                                                                                                                         |
|-------------------------|-------------------------------------------------------------------------------------------------------------------------|
| <b>Description</b>      | Unconjugated Rabbit polyclonal to PRD10                                                                                 |
| <b>Model</b>            | STJ191198                                                                                                               |
| <b>Host</b>             | Rabbit                                                                                                                  |
| <b>Reactivity</b>       | Human, Mouse                                                                                                            |
| <b>Applications</b>     | ELISA, WB                                                                                                               |
| <b>Immunogen</b>        | Synthesized peptide derived from human PRD10 protein.                                                                   |
| <b>Immunogen Region</b> | 280-360aa                                                                                                               |
| <b>Gene ID</b>          | <a href="#">56980</a>                                                                                                   |
| <b>Gene Symbol</b>      | <a href="#">PRDM10</a>                                                                                                  |
| <b>Dilution range</b>   | WB 1:500-2000 ELISA 1:5000-20000                                                                                        |
| <b>Specificity</b>      | PRD10 Polyclonal Antibody detects endogenous levels of protein.                                                         |
| <b>Purification</b>     | PRD10 antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen. |
| <b>Note</b>             | For Research Use Only (RUO).                                                                                            |
| <b>Protein Name</b>     | PR domain zinc finger protein 10 PR domain-containing protein 10 Tristanin                                              |
| <b>Molecular Weight</b> | 126 kDa                                                                                                                 |
| <b>Clonality</b>        | Polyclonal                                                                                                              |
| <b>Conjugation</b>      | Unconjugated                                                                                                            |
| <b>Isotype</b>          | IgG                                                                                                                     |

|                            |                                                                            |
|----------------------------|----------------------------------------------------------------------------|
| <b>Formulation</b>         | Liquid form in PBS containing 50% glycerol, and 0.02% sodium azide.        |
| <b>Concentration</b>       | 1 mg/ml                                                                    |
| <b>Storage Instruction</b> | Store at -20°C, and avoid repeat freeze-thaw cycles.                       |
| <b>Database Links</b>      | <a href="#">HGNC:13995</a> OMIM:NA                                         |
| <b>Alternative Names</b>   | PR domain zinc finger protein 10 PR domain-containing protein 10 Tristanin |

---

**St John's Laboratory Ltd**

**F** +44 (0)207 681 2580

**T** +44 (0)208 223 3081

**W** <http://www.stjohnslabs.com/>

**E** [info@stjohnslabs.com](mailto:info@stjohnslabs.com)