

Anti-RAB17 antibody



Description Unconjugated Rabbit polyclonal to RAB17

Model STJ191271

Host Rabbit

Reactivity Human, Mouse

Applications ELISA, WB

Immunogen Synthesized peptide derived from human RAB17 protein.

Immunogen Region 50-130aa

Gene ID <u>64284</u>

Gene Symbol RAB17

Dilution range WB 1:500-2000 ELISA 1:5000-20000

Specificity RAB17 Polyclonal Antibody detects endogenous levels of protein.

Tissue Specificity Expressed in melanocytes (at protein level).

Purification RAB17 antibody was affinity-purified from rabbit antiserum by affinity-

chromatography using epitope-specific immunogen.

Note For Research Use Only (RUO).

Protein Name Ras-related protein Rab-17

Molecular Weight 23 kDa

Clonality Polyclonal

Conjugation Unconjugated

Isotype IgG

Formulation Liquid form in PBS containing 50% glycerol, and 0.02% sodium azide.

Concentration 1 mg/ml

Storage Instruction Store at -20°C, and avoid repeat freeze-thaw cycles.

Database Links HGNC:16523OMIM:602206

Alternative Names Ras-related protein Rab-17

Function The small GTPases Rab are key regulators of intracellular membrane

trafficking, from the formation of transport vesicles to their fusion with membranes. Rabs cycle between an inactive GDP-bound form and an active GTP-bound form that is able to recruit to membranes different set of downstream effectors directly responsible for vesicle formation, movement, tethering and fusion. That Rab is involved in transcytosis, the directed movement of endocytosed material through the cell and its executosis from

movement of endocytosed material through the cell and its exocytosis from the plasma membrane at the opposite side. Mainly observed in epithelial cells, transcytosis mediates for instance, the transcellular transport of

immunoglobulins from the basolateral surface to the apical surface. Most probably controls membrane trafficking through apical recycling endosomes in a post-endocytic step of transcytosis. Required for melanosome transport and release from melanocytes, it also regulates dendrite and dendritic spine

development. May also play a role in cell migration.

Cellular Localization Recycling endosome membrane Melanosome Cell projection, dendrite. May

also localize at the basolateral and apical plasma membrane. In neurons,

localizes to the cell body and dendritic shaft and spine.

St John's Laboratory Ltd

F +44 (0)207 681 2580

T +44 (0)208 223 3081

W http://www.stjohnslabs.com/ E info@stjohnslabs.com