

Anti-SCN3A antibody



Description Unconjugated Rabbit polyclonal to SCN3A

Model STJ191436

Host Rabbit

Reactivity Human

Applications ELISA, WB

Gene ID <u>6328</u>

Gene Symbol SCN3A

Dilution range WB 1:500-2000 ELISA 1:5000-20000

Specificity SCN3A Polyclonal Antibody detects endogenous levels of protein.

Purification SCN3A antibody was affinity-purified from rabbit antiserum by affinity-

chromatography using epitope-specific immunogen.

Note For Research Use Only (RUO).

Protein Name Sodium channel protein type 3 subunit alpha Sodium channel protein brain III

subunit alpha Sodium channel protein type III subunit alpha Voltage-gated sodium channel subtype III Voltage-gated sodium channel subunit alpha

Nav1.3

Molecular Weight 220 kDa

Clonality Polyclonal

Conjugation Unconjugated

Isotype IgG

Formulation Liquid form in PBS containing 50% glycerol, and 0.02% sodium azide.

Concentration 1 mg/ml

Storage Instruction Store at -20°C, and avoid repeat freeze-thaw cycles.

Database Links <u>HGNC:10590OMIM:182391</u>

Alternative Names Sodium channel protein type 3 subunit alpha Sodium channel protein brain III

subunit alpha Sodium channel protein type III subunit alpha Voltage-gated sodium channel subtype III Voltage-gated sodium channel subunit alpha

Nav1.3

Function Mediates the voltage-dependent sodium ion permeability of excitable

membranes. Assuming opened or closed conformations in response to the voltage difference across the membrane, the protein forms a sodium-selective channel through which Na(+) ions may pass in accordance with their

electrochemical gradient.

Sequence and Domain Family The sequence contains 4 internal repeats, each with 5 hydrophobic segments

(S1, S2, S3, S5, S6) and one positively charged segment (S4). Segments S4 are probably the voltage-sensors and are characterized by a series of positively

charged amino acids at every third position.

Cellular Localization Cell membrane

Post-translational May be ubiquitinated by NEDD4L; which would promote its endocytosis.; **Modifications** Phosphorylation at Ser-1501 by PKC in a highly conserved cytoplasmic loc

Phosphorylation at Ser-1501 by PKC in a highly conserved cytoplasmic loop slows inactivation of the sodium channel and reduces peak sodium currents.

St John's Laboratory Ltd

F +44 (0)207 681 2580

T+44 (0)208 223 3081

W http://www.stjohnslabs.com/ E info@stjohnslabs.com