

Anti-UBP14 antibody



Description	Unconjugated Rabbit polyclonal to UBP14
Model	STJ191570
Host	Rabbit
Reactivity	Human, Mouse
Applications	ELISA, WB
Gene ID	9097
Gene Symbol	USP14
Dilution range	WB 1:500-2000 ELISA 1:5000-20000
Specificity	UBP14 Polyclonal Antibody detects endogenous levels of protein.
Purification	UBP14 antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
Note	For Research Use Only (RUO).
Protein Name	Ubiquitin carboxyl-terminal hydrolase 14 Deubiquitinating enzyme 14 Ubiquitin thioesterase 14 Ubiquitin-specific-processing protease 14
Molecular Weight	54 kDa
Clonality	Polyclonal
Conjugation	Unconjugated
Isotype	IgG
Formulation	Liquid form in PBS containing 50% glycerol, and 0.02% sodium azide.

Concentration	1 mg/ml
Storage Instruction	Store at -20°C, and avoid repeat freeze-thaw cycles.
Database Links	HGNC:12612OMIM:607274
Alternative Names	Ubiquitin carboxyl-terminal hydrolase 14 Deubiquitinating enzyme 14 Ubiquitin thioesterase 14 Ubiquitin-specific-processing protease 14
Function	Proteasome-associated deubiquitinase which releases ubiquitin from the proteasome targeted ubiquitinated proteins. Ensures the regeneration of ubiquitin at the proteasome. Is a reversibly associated subunit of the proteasome and a large fraction of proteasome-free protein exists within the cell. Required for the degradation of the chemokine receptor CXCR4 which is critical for CXCL12-induced cell chemotaxis. Serves also as a physiological inhibitor of endoplasmic reticulum-associated degradation (ERAD) under the non-stressed condition by inhibiting the degradation of unfolded endoplasmic reticulum proteins via interaction with ERN1. Indispensable for synaptic development and function at neuromuscular junctions (NMJs).
Cellular Localization	Cytoplasm Cell membrane

St John's Laboratory Ltd

F +44 (0)207 681 2580

T +44 (0)208 223 3081

W <http://www.stjohnslabs.com/>

E info@stjohnslabs.com