

## Anti-CCNK antibody

---



<b>Description</b>	Unconjugated Rabbit polyclonal to CCNK
<b>Model</b>	STJ191667
<b>Host</b>	Rabbit
<b>Reactivity</b>	Human, Mouse
<b>Applications</b>	ELISA, WB
<b>Immunogen</b>	Synthesized peptide derived from human CCNK protein.
<b>Immunogen Region</b>	80-160aa
<b>Gene ID</b>	<a href="#">8812</a>
<b>Gene Symbol</b>	<a href="#">CCNK</a>
<b>Dilution range</b>	WB 1:500-2000 ELISA 1:5000-20000
<b>Specificity</b>	CCNK Polyclonal Antibody detects endogenous levels of protein.
<b>Tissue Specificity</b>	Ubiquitously expressed. Highest levels in testis.
<b>Purification</b>	CCNK antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
<b>Note</b>	For Research Use Only (RUO).
<b>Protein Name</b>	Cyclin-K
<b>Molecular Weight</b>	63 kDa
<b>Clonality</b>	Polyclonal
<b>Conjugation</b>	Unconjugated

<b>Isotype</b>	IgG
<b>Formulation</b>	Liquid form in PBS containing 50% glycerol, and 0.02% sodium azide.
<b>Concentration</b>	1 mg/ml
<b>Storage Instruction</b>	Store at -20°C, and avoid repeat freeze-thaw cycles.
<b>Database Links</b>	<a href="#">HGNC:1596</a> <a href="#">OMIM:603544</a>
<b>Alternative Names</b>	Cyclin-K
<b>Function</b>	Regulatory subunit of cyclin-dependent kinases that mediates activation of target kinases. Plays a role in transcriptional regulation via its role in regulating the phosphorylation of the C-terminal domain (CTD) of the large subunit of RNA polymerase II (POLR2A).
<b>Cellular Localization</b>	Nucleus

---

**St John's Laboratory Ltd**

**F** +44 (0)207 681 2580

**T** +44 (0)208 223 3081

**W** <http://www.stjohnslabs.com/>

**E** [info@stjohnslabs.com](mailto:info@stjohnslabs.com)