

## Anti-CCNL2 antibody

---



<b>Description</b>	Unconjugated Rabbit polyclonal to CCNL2
<b>Model</b>	STJ191668
<b>Host</b>	Rabbit
<b>Reactivity</b>	Human, Mouse, Rat
<b>Applications</b>	ELISA, WB
<b>Immunogen</b>	Synthesized peptide derived from human CCNL2 protein.
<b>Immunogen Region</b>	400-480aa
<b>Gene ID</b>	<a href="#">81669</a>
<b>Gene Symbol</b>	<a href="#">CCNL2</a>
<b>Dilution range</b>	WB 1:500-2000 ELISA 1:5000-20000
<b>Specificity</b>	CCNL2 Polyclonal Antibody detects endogenous levels of protein.
<b>Tissue Specificity</b>	Widely expressed.
<b>Purification</b>	CCNL2 antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
<b>Note</b>	For Research Use Only (RUO).
<b>Protein Name</b>	Cyclin-L2 Paneth cell-enhanced expression protein
<b>Molecular Weight</b>	57 kDa
<b>Clonality</b>	Polyclonal
<b>Conjugation</b>	Unconjugated

<b>Isotype</b>	IgG
<b>Formulation</b>	Liquid form in PBS containing 50% glycerol, and 0.02% sodium azide.
<b>Concentration</b>	1 mg/ml
<b>Storage Instruction</b>	Store at -20°C, and avoid repeat freeze-thaw cycles.
<b>Database Links</b>	<a href="https://www.ncbi.nlm.nih.gov/condensedcode/HGNC:20570OMIM:613482">HGNC:20570OMIM:613482</a>
<b>Alternative Names</b>	Cyclin-L2 Paneth cell-enhanced expression protein
<b>Function</b>	Involved in pre-mRNA splicing. May induce cell death, possibly by acting on the transcription and RNA processing of apoptosis-related factors.
<b>Sequence and Domain Family</b>	Contains a RS region (arginine-serine dipeptide repeat) within the C-terminal domain which is the hallmark of the SR family of splicing factors. This region probably plays a role in protein-protein interactions .
<b>Cellular Localization</b>	Nucleus speckle Nucleus, nucleoplasm

---

**St John's Laboratory Ltd**

**F** +44 (0)207 681 2580

**T** +44 (0)208 223 3081

**W** <http://www.stjohnslabs.com/>

**E** [info@stjohnslabs.com](mailto:info@stjohnslabs.com)