

Anti-CCNL2 antibody



Description	Unconjugated Rabbit polyclonal to CCNL2
Model	STJ191668
Host	Rabbit
Reactivity	Human, Mouse, Rat
Applications	ELISA, WB
Immunogen	Synthesized peptide derived from human CCNL2 protein.
Immunogen Region	400-480aa
Gene ID	81669
Gene Symbol	CCNL2
Dilution range	WB 1:500-2000 ELISA 1:5000-20000
Specificity	CCNL2 Polyclonal Antibody detects endogenous levels of protein.
Tissue Specificity	Widely expressed.
Purification	CCNL2 antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
Note	For Research Use Only (RUO).
Protein Name	Cyclin-L2 Paneth cell-enhanced expression protein
Molecular Weight	57 kDa
Clonality	Polyclonal
Conjugation	Unconjugated

Isotype	IgG
Formulation	Liquid form in PBS containing 50% glycerol, and 0.02% sodium azide.
Concentration	1 mg/ml
Storage Instruction	Store at -20°C, and avoid repeat freeze-thaw cycles.
Database Links	HGNC:20570OMIM:613482
Alternative Names	Cyclin-L2 Paneth cell-enhanced expression protein
Function	Involved in pre-mRNA splicing. May induce cell death, possibly by acting on the transcription and RNA processing of apoptosis-related factors.
Sequence and Domain Family	Contains a RS region (arginine-serine dipeptide repeat) within the C-terminal domain which is the hallmark of the SR family of splicing factors. This region probably plays a role in protein-protein interactions .
Cellular Localization	Nucleus speckle Nucleus, nucleoplasm

St John's Laboratory Ltd

F +44 (0)207 681 2580

T +44 (0)208 223 3081

W <http://www.stjohnslabs.com/>

E info@stjohnslabs.com