

Anti-CD27 antibody



Description	Unconjugated Rabbit polyclonal to CD27
Model	STJ191672
Host	Rabbit
Reactivity	Human
Applications	ELISA, WB
Immunogen	Synthesized peptide derived from human CD27 protein.
Immunogen Region	1-80aa
Gene ID	939
Gene Symbol	CD27
Dilution range	WB 1:500-2000 ELISA 1:5000-20000
Specificity	CD27 Polyclonal Antibody detects endogenous levels of protein.
Tissue Specificity	Found in most T-lymphocytes.
Purification	CD27 antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
Note	For Research Use Only (RUO).
Protein Name	CD27 antigen CD27L receptor T-cell activation antigen CD27 T14 Tumor necrosis factor receptor superfamily member 7 CD antigen CD27
Molecular Weight	28 kDa
Clonality	Polyclonal

Conjugation	Unconjugated
Isotype	IgG
Formulation	Liquid form in PBS containing 50% glycerol, and 0.02% sodium azide.
Concentration	1 mg/ml
Storage Instruction	Store at -20°C, and avoid repeat freeze-thaw cycles.
Database Links	HGNC:11922OMIM:186711
Alternative Names	CD27 antigen CD27L receptor T-cell activation antigen CD27 T14 Tumor necrosis factor receptor superfamily member 7 CD antigen CD27
Function	Receptor for CD70/CD27L. May play a role in survival of activated T-cells. May play a role in apoptosis through association with SIVA1.
Cellular Localization	Membrane. Single-pass type I membrane protein.
Post-translational Modifications	Phosphorylated.; O-glycosylated with core 1 or possibly core 8 glycans.

St John's Laboratory Ltd

F +44 (0)207 681 2580

T +44 (0)208 223 3081

W <http://www.stjohnslabs.com/>

E info@stjohnslabs.com