

Anti-SNPC3 antibody



| | |
|-------------------------|--|
| Description | Unconjugated Rabbit polyclonal to SNPC3 |
| Model | STJ191777 |
| Host | Rabbit |
| Reactivity | Human |
| Applications | ELISA, WB |
| Gene ID | 6619 |
| Gene Symbol | SNAPC3 |
| Dilution range | WB 1:500-2000 ELISA 1:5000-20000 |
| Specificity | SNPC3 Polyclonal Antibody detects endogenous levels of protein. |
| Purification | SNPC3 antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen. |
| Note | For Research Use Only (RUO). |
| Protein Name | snRNA-activating protein complex subunit 3 SNAPc subunit 3 Proximal sequence element-binding transcription factor subunit beta PSE-binding factor subunit beta PTF subunit beta Small nuclear RNA-activating complex polypeptide |
| Molecular Weight | 45 kDa |
| Clonality | Polyclonal |
| Conjugation | Unconjugated |
| Isotype | IgG |

| | |
|------------------------------|---|
| Formulation | Liquid form in PBS containing 50% glycerol, and 0.02% sodium azide. |
| Concentration | 1 mg/ml |
| Storage Instruction | Store at -20°C, and avoid repeat freeze-thaw cycles. |
| Database Links | HGNC:11136OMIM:602348 |
| Alternative Names | snRNA-activating protein complex subunit 3 SNAPc subunit 3 Proximal sequence element-binding transcription factor subunit beta PSE-binding factor subunit beta PTF subunit beta Small nuclear RNA-activating complex polypeptide |
| Function | Part of the SNAPc complex required for the transcription of both RNA polymerase II and III small-nuclear RNA genes. Binds to the proximal sequence element (PSE), a non-TATA-box basal promoter element common to these 2 types of genes. Recruits TBP and BRF2 to the U6 snRNA TATA box. |
| Cellular Localization | Nucleus. |

St John's Laboratory Ltd

F +44 (0)207 681 2580

T +44 (0)208 223 3081

W <http://www.stjohnslabs.com/>

E info@stjohnslabs.com