

Anti-DEDD antibody



Description Unconjugated Rabbit polyclonal to DEDD

Model STJ191839

Host Rabbit

Reactivity Human, Mouse, Rat

Applications ELISA, WB

Immunogen Synthesized peptide derived from human DEDD protein.

Immunogen Region 90-170aa

Gene ID 9191

Gene Symbol <u>DEDD</u>

Dilution range WB 1:500-2000 ELISA 1:5000-20000

Specificity DEDD Polyclonal Antibody detects endogenous levels of protein.

Tissue Specificity Widely expressed with highest levels in testis.

Purification DEDD antibody was affinity-purified from rabbit antiserum by affinity-

chromatography using epitope-specific immunogen.

Note For Research Use Only (RUO).

Protein Name Death effector domain-containing protein DEDPro1 Death effector domain-

containing testicular molecule FLDED-1

Molecular Weight 34 kDa

Clonality Polyclonal

Conjugation Unconjugated

Isotype IgG

Formulation Liquid form in PBS containing 50% glycerol, and 0.02% sodium azide.

Concentration 1 mg/ml

Storage Instruction Store at -20°C, and avoid repeat freeze-thaw cycles.

Database Links <u>HGNC:27550MIM:606841</u>

Alternative Names Death effector domain-containing protein DEDPro1 Death effector domain-

containing testicular molecule FLDED-1

Function A scaffold protein that directs CASP3 to certain substrates and facilitates their

ordered degradation during apoptosis. May also play a role in mediating CASP3 cleavage of KRT18. Regulates degradation of intermediate filaments during apoptosis. May play a role in the general transcription machinery in the nucleus and might be an important regulator of the activity of GTF3C3.

Inhibits DNA transcription in vitro .

Cytoplasm. Nucleus, nucleolus. Translocated to the nucleus during CD95-

mediated apoptosis where it is localized in the nucleoli . Following apoptosis induction, the mono and/or diubiquitination form increases and forms filamentous structures that colocalize with KRT8 and KRT18 intermediate

filament network in simple epithelial cells.

Post-translational

Modifications

Exists predominantly in a mono- or diubiquitinated form.

St John's Laboratory Ltd

F +44 (0)207 681 2580

T+44 (0)208 223 3081

W http://www.stjohnslabs.com/ E info@stjohnslabs.com